

Research article

## Citrus Flavonoid Supplementation Improves Exercise Performance in Trained Athletes

Elvera Overdevest<sup>1</sup>, Jeroen A. Wouters<sup>1</sup>, Kevin H.M. Wolfs<sup>2</sup>, Job J.M. van Leeuwen<sup>2</sup> and Sam Possemiers<sup>2</sup>✉

<sup>1</sup>Team NL Innovation Center, Sportcentrum Papendal, Arnhem, The Netherlands; <sup>2</sup>BioActor BV, Maastricht, The Netherlands

### Abstract

Previous studies have shown that polyphenol supplementation may be an effective strategy to improve exercise performance, due to their antioxidant character and ability to stimulate NO production. These properties may contribute to exercise performance, yet no conclusive research has been performed in exploring the direct effects of citrus flavonoids on human exercise performance. Therefore, the purpose of this study was to assess whether supplementation of a customized citrus flavonoid (CF) extract for 4 weeks improves cycling time-trial performance in trained male athletes. In a double-blind, randomized, parallel study, 39 healthy, trained males were given a daily dose of either 500 mg of a customized citrus flavonoid extract (CF) or a placebo for 4 weeks. Exercise performance was tested by means of a time-trial test on a cycle ergometer, during which participants had to generate as much power as possible for duration of 10 minutes. Absolute power output significantly increased with  $14.9 \pm 3.9$  W after 4 weeks of CF supplementation, corresponding with a 5.0% increase, compared to  $3.8 \pm 3.2$  W (1.3% increase) in placebo ( $p < 0.05$ ). In addition, oxygen consumption/power ratio significantly decreased in the CF group compared to placebo ( $p = 0.001$ ), and a trend was found in the change in peak power output in CF ( $18.2 \pm 23.2$  W) versus placebo ( $-28.4 \pm 17.6$  W;  $p = 0.116$ ). The current study is the first convincing report that citrus flavonoid supplementation can improve exercise performance, as shown by a significant increase in power output during the exercise test.

**Key words:** Hesperetin, power output, antioxidant, time trial, ergometer.

### Introduction

Sport nutrition and nutritional supplementation aimed at enhancing exercise performance and recovery from exercise is commonly used by athletes (Burke and Read, 1993; Froiland et al., 2004; Tscholl et al., 2010). For instance, foods high in carbohydrates and protein are consumed in order to replenish glycogen stores and promote muscle anabolism (Reid, 2013). Although exercise is known to be beneficial for improving exercise performance and preventing several pathological conditions, like cardiovascular disease, type II diabetes, metabolic syndrome, and neurodegenerative diseases, it also induces reactive oxygen species (ROS) production, which is related with several conditions leading to diseases (Davis et al., 2010; Davis et al., 2009; Halliwell, 1991; Malaguti et al., 2013; Masella et al., 2005). Exhaustive exercise increases

ROS production, leading to muscle fiber damage, which eventually results in muscle fatigue (Peternej and Coombes, 2011). However, there is growing evidence suggesting that the presence of a small stimulus, like low concentration of ROS, is able to express the transcription of major antioxidant genes. Enzymes like superoxide dismutase (SOD) and glutathione are important antioxidant defenses that protect cells from ROS-induced oxidative stress (Masella et al., 2005). Moderate exercise acts as a stimulator of the body's antioxidant defenses against oxidative damage (Gomez-Cabrera et al., 2008; Powers et al., 2011). The correlation between oxidative damage and muscle fatigue could be an important strategy for nutritional interventions to increase exercise performance. Antioxidant supplementation may be an effective strategy, considering the reactive oxygen species (ROS) scavenging effects that could lead to a reduction in muscle damage caused by prolonged exercise (Myburgh, 2014; Sachdev and Davies, 2008).

Polyphenols, including flavonoids derived primarily from fruits, have been of interest due to their antioxidant and anti-inflammatory effects (Masella et al., 2005). Previous studies showed that polyphenols derived from pomegranates, cherries and blueberries reduced muscle soreness and improved muscle strength after eccentric exercise (Bowtell et al., 2011; McLeay et al., 2012; Trombold et al., 2010; 2011). Next to potential benefits in muscle recovery, flavonoid supplementation has been shown to improve endurance exercise performance in humans (Davis et al., 2009; 2010; MacRae and Mefferd, 2006; Nieman et al., 2010).

Another point of interest is the association that has been made with stimulation of nitric oxide (NO) production and subsequent improved endothelial function with different flavonoids, including hesperidin (Rizza et al., 2011). NO acts on smooth muscle cells within the arterial wall, causing vasodilatation and thereby improving blood flow and reducing blood pressure (Cooke et al., 1997; Umans and Levi, 1995). During exercise, this allows for an increase in transport of oxygen and nutrients to the muscles and removes metabolites causing fatigue in prolonged exercise (Bescós et al., 2012; Bloomer, 2010; Wylie et al., 2013).

However, no conclusive clinical research has been performed to explore direct benefits of citrus flavonoids on exercise performance in trained athletes (Malaguti et al., 2013). Therefore, the aim of the study was to elucidate whether a customized citrus flavonoid (CF) extract sup-

plementation for duration of 4 weeks improves cycling time-trial performance in trained male athletes. It was hypothesized that CF extract supplementation will improve time-trial performance in trained athletes.

## Methods

### Participants

Trained, non-smoking male participants, aged 18 – 25 y, engaging in moderate to high physical activity for a minimum of 30 minutes at least 3 times a week were included in the study. Exclusion criteria were the use of dietary antioxidant and/or vitamin supplements, and the ingestion of products containing citrus flavonoids or its metabolite 4 days prior to and during the study.

All participants gave written informed consent before participation. The study was approved by the Medical Ethics Committee of Wageningen University and conducted in full accordance with the principles of the Declaration of Helsinki of 1975 as amended in 2013 and with the Dutch Regulations on Medical Research involving Human Subjects (WMO, 1998). The study was performed at InnoSportlab Papendal, Arnhem, The Netherlands (now part of the National Sportscenter Papendal). The trial has been registered in the Clinical Trials register (NCT02787733).

### Design and protocol

The study was designed as a randomized, parallel group, double-blind design to test the effects of a daily dose of 500 mg CF extract supplementation versus a placebo over a period of 4 weeks. A publicly available randomization program (<http://randomizer.org>) was used to randomly assign participants to one of the two interventions.

The study consisted of a familiarization test, a baseline test (Test 1) and a final test after a 4-week intervention period (Test 2). Prior to each test, participants were instructed to standardize their workouts and dietary intake and to refrain from physical exercise and alcohol for at least 24-h prior to testing. During the complete duration of the study, participants were instructed to refrain from eating foods containing citrus flavonoids, including lemons, oranges, and grapefruit.

The familiarization test consisted of a 5 min warm-up at 100 W followed by a 10 min time-trial on a cycle ergometer (SRM, Jülich, Germany), during which participants had to generate the maximal power (W) possible over the time course of 10 min. Power output (W) and heart rate (bpm) were measured continuously during the time-trial, and indirect calorimetry was used to measure oxygen consumption ( $\text{VO}_2$ ;  $\text{mL}\cdot\text{min}^{-1}$ ) during the time-trial. In addition, an estimation of maximal oxygen consumption ( $\text{VO}_{2\text{max}}$ ;  $\text{mL}\cdot\text{min}^{-1}$ ) was made. At  $t = 0, 9, 10$  and  $11$  min after starting the test, participants were asked to indicate their perceived exhaustion (RPE) using a Borg scale from 6 – 20, in which 6 is no exertion at all and 20 is maximal exertion. The mean power that was measured during the familiarization test was used to determine the power for a 10 min pre-exhaustion test during Test 1 and Test 2.

The following test protocol was performed identi-

cally on both test days. First, participants had to perform a standardized pre-exhaustion test, consisting of 10 min cycling at 80% of the mean work-load as established during the familiarization test. During this pre-exhaustion phase, heart rate and exhaustion at  $t = 0, 9, 10$  and  $11$  min after start of the test were measured. Subsequently, participants took a passive rest period of 25 min followed by 5 min of warm-up at 100 W, as a warm-up has been shown to improve subsequent exercise performance (Fradkin et al., 2010). Directly after that, participants had to perform a 10 min time-trial comparable to the familiarization test. The same parameters were obtained as in the familiarization test with additional exhaustion measurements at  $t = -2, -1$  prior to start of the test. Between Test 1 and Test 2, an intervention period of 4 weeks was conducted, in which the participants administered either CF extract supplementation or placebo supplement.

### Study product

BioActor BV (Maastricht, the Netherlands) supplied the CF extract, standardized for hesperetin-7-O-rutinoside 2S enantiomer, with a total rutinoid content of at least 90% (WATTS'UP<sup>®</sup>/CF extract). The study product was formulated into capsules containing 250 mg of CF extract and 10 mg Magnesium Stearate by Aminolabs (Hasselt, Belgium). The placebo capsules containing 250 mg Microcrystalline Cellulose were produced to be identical in appearance and taste. The capsules were transported and stored at room temperature. Participants were instructed to ingest two capsules with 200 mL water daily before breakfast, for 4 weeks from the first morning after Test 1 (T1) until the morning of Test 2 (T2).

### Statistical analysis

Statistical analysis was performed using IBM SPSS Statistics for Windows (version 22.0 Armonk, NY, USA). Only data from participants who completed the study ( $n = 39$ ) were used in the final analysis. Before statistical analysis, the Shapiro-Wilk test was performed to test for normality in the test population. To determine the effects of CF supplementation on the measured performance outcomes, the mean values at baseline (T1) were compared with the mean values after 4 weeks of supplementation (T2). A paired-samples t-test was used to compare the T1 and T2 measurements within both the CF and placebo group. To compare the differences between the CF and placebo group, a two-way mixed ANOVA was performed. Data is reported as mean  $\pm$  standard error of the mean (SEM) and the level of significance was set on  $p < 0.05$ .

## Results

### Study participants

Thirty-nine participants completed the study protocol, as one participant dropped out after the first test day due to an injury not related to the study. The baseline characteristics of the participants in both groups are shown in Table 1. No significant differences were found between CF extract supplementation and placebo at the start of the 4 week intervention period.

**Table 1.** Participant characteristics of the 39 participants that completed the study protocol. Data were obtained at baseline and values are expressed as mean ( $\pm$  SEM).

	Total (n=39)	Citrus Flavonoid (CF) (n=19)	Placebo (n=20)
Age (yr)	23.0 (.3)	22.7 (.4)	23.6 (.4)
Weight (kg)	74.4 (1.29)	74.1 (1.75)	74.8 (1.93)
Height (m)	1.84 (.01)	1.83 (.02)	1.84 (.02)
BMI (kg·m <sup>-2</sup> )	22.1 (.30)	22.1 (.42)	22.0 (.44)
Exercise time per week	9.6 (.6)	8.8 (.6)	10.3 (.9)
Absolute power (W)	299.3 (7.7)	298.0 (10.8)	300.5 (11.4)
Relative power (W·kg <sup>-1</sup> )	4.0 (.1)	4.0 (.2)	4.0 (.2)
Heart rate (BPM)	174 (2)	171 (4)	177 (3)
Average VO <sub>2</sub> (mL·kg <sup>-1</sup> ·min)	50.3 (1.3)	51.1 (1.6)	49.6 (2.1)
Estimated VO <sub>2max</sub> (mL·kg <sup>-1</sup> ·min)	57.1 (1.2)	57.9 (1.6)	56.5 (1.8)
VO <sub>2</sub> /Power ratio	.172 (.003)	.176 (.005)	.170 (.004)
Peak power output (W)	332.5 (21.0)	359.4 (36.0)	306.8 (22.0)

No significant differences were found between CF and placebo group at baseline,  $p > 0.05$ .

**Table 2.** Within and between group comparison of the performance outcomes of the citrus flavonoid group and placebo group, with corresponding p-values. Values are expressed as mean ( $\pm$  SEM).


Variables	Citrus Flavonoid (CF) (n=19)				Placebo (n=20)				
	Test 1	Test 2	$\Delta$ T2-T1	<P1†	Test 1	Test 2	$\Delta$ T2-T1	<P2‡	<P3§
Weight (kg)	74.1 (1.8)	74.2 (1.74)	.1 (1)	.406	74.8 (1.9)	74.7 (1.9)	-.1 (.4)	.718	.529
Abs P (W)	298 (11)	313 (10)	14.9 (3.9)	.001	301 (11)	304.3 (11.4)	3.8 (3.2)	.243	.032
Rel P (W·kg <sup>-1</sup> )	4.0 (.1)	4.2 (.1)	.2 (1)	.002	4.0 (.2)	4.1 (1)	.1 (1)	.328	.077
HR (BPM)	170 (4)	169 (3)	-1 (2)	.578	176 (3)	177 (3)	1 (1)	.206	.276
Av VO <sub>2</sub>	51.1 (1.6)	52.0 (1.2)	.9 (7)	.235	49.6 (2.1)	50.2 (1.8)	.6 (.6)	.330	.736
Es VO <sub>2max</sub>	57.9 (1.6)	57.9 ( $\pm$ 1.3)	.0 (7)	.988	56.5 (1.8)	56.7 (1.6)	.2 (.9)	.784	.826
VO <sub>2</sub> /Power ratio	.176 (.005)	.167 (.004)	-.009 (.002)	.001	.170 (.004)	.171 (.004)	.001 (.002)	.534	.001
PP output (W)	360 (36)	378 (40)	18 (23)	.444	307 (22)	279 (20)	-29 (17)	.123	.116


Abs P: Absolute power; Rel P: Relative power; HR: Heart rate; Av VO<sub>2</sub>: Average VO<sub>2</sub> (mL·kg<sup>-1</sup>·min<sup>-1</sup>); Es VO<sub>2max</sub>: Estimated VO<sub>2max</sub> (mL·kg<sup>-1</sup>·min<sup>-1</sup>); PP output: Peak power output. †P1; P-value of within CF group differences between Test 2 and Test 1 ( $\Delta$ T2-T1), compared using a paired t-test, ‡P2; P-value of within placebo group differences between Test 2 and Test 1 ( $\Delta$ T2-T1), compared using a paired t-test, §P3; P-value of the difference between pre- and post-treatment between the CF and placebo group using a mixed ANOVA.

### Exercise performance, power output, and VO<sub>2</sub>

Table 2 shows the results of the performance outcomes from the participants with CF extract supplementation and placebo, before and after the 4 week intervention period. Both absolute ( $p = 0.001$ ) and relative ( $p = 0.002$ ) power output were significantly increased over the 4 week supplementation in the group receiving CF extract supplementation. No difference in weight was noted during the trial period in either group, making that the relative power parameter in this study did not need to be corrected for changes in weight. In the placebo group, no significant differences in absolute and relative power were found when comparing T2 with T1. The power output of each participant for the whole duration of the time-trial test was also recorded. Figure 1 shows the average power evolution profiles for both groups at baseline and after 4 weeks of CF extract supplementation.

When plotting the mean absolute power output (Figure 2A), it becomes apparent that the group receiving CF extract supplementation significantly increased ( $p = 0.001$ ) absolute power output by 5% over the 4-week supplementation period, while the placebo group did not ( $p > 0.05$ ). When expressed as the difference in absolute power output between test days ( $\Delta$ T2 - T1), the increase in power after 4 weeks of CF extract supplementation is significant and almost 4 times higher than placebo ( $p = 0.032$ ; Figure 2B). In addition, a trend was found for the increase in peak power output (in the first 3 seconds of workout) after 4 weeks of treatment in CF extract supplementation ( $18.2 \pm 23.2$  W) versus placebo ( $-29.0 \pm 17.4$  W;  $p = 0.116$ ; Table 2).

**Figure 1.** A) Average power evolution profiles during Test 1 for CF and placebo group. B) Average power evolution profiles during Test 2 for CF and placebo group.



**Figure 2.** A) After 4 weeks of supplementation, the average power in the CF group was increased with 5% ( $p = 0.001^{**}$ ), while no significant increase was noted in the placebo group ( $p = 0.243$ ). Values are expressed as mean  $\pm$  SEM. B) The increase in power after 4 weeks treatment compared to baseline ( $\Delta T_2 - T_1$ ) is significantly higher within the CF group than within the placebo group ( $p = 0.032^*$ ). Values are expressed as mean  $\pm$  SEM.


In addition to power output, the  $VO_2$  consumption was measured. No significant differences ( $p > 0.05$ ) in  $VO_2$  consumption were found between baseline and after four weeks of treatments in both groups (Figure 3A). When plotting the  $VO_2$  consumption against the power output data, the results show that the  $VO_2$ /Power ratio significantly decreased ( $p = 0.001$ ) with 5.1% in the CF extract supplementation group compared to a non-significant 0.6% increase ( $p = 0.534$ ) in the placebo group (Figure 3B). When looking at the delta  $VO_2$ /Power ratio ( $\Delta T_2 - T_1$ ), a significant between-group effect is apparent ( $p = 0.001$ ), in which the  $VO_2$ /Power ratio decreased with 0.009 in the CF extract supplementation group and insignificantly increased with 0.001 in the placebo group.

## Discussion

The present study shows that CF extract supplementation for 4 weeks significantly increased absolute power in trained athletes compared to placebo. Furthermore, body mass and  $VO_2$  did not change in either groups, resulting in a higher amount of power produced per kilogram of bodyweight (relative power) and per unit of oxygen ( $VO_2$ /Power ratio) in the CF group compared to placebo.

Previous studies showed that the flavonoid quercetin increased exercise performance and endurance capacity (Davis et al., 2010; MacRae and Mefferd, 2006). While

*in vivo* the exact mode of action of flavonoids is unclear, oxygraphic results for quercetin and hesperetin support the shared working mechanism of both compounds by which they can increase exercise performance (Dorta et al., 2005).



**Figure 3.** A) Average  $VO_2$  consumption over time in both groups, before and after 4 weeks of intervention. B) The  $VO_2$ /power ratio was significantly decreased in the CF group compared to the ratio at baseline ( $p = 0.001^{**}$ ). Values are expressed as mean  $\pm$  SEM.

In addition to their antioxidant activity, flavonoids may exert other intracellular effects and may interact at the mitochondrial level, contributing to maintain high energetic demand and redox homeostasis. It is suggested that polyphenols may have a potential effect on the functioning of the whole electron transfer chain (ETC), acting on the total mitochondrial respiration process rather than focusing on only a single specific complex (e.g. Complex I) (Santos et al., 1998). In addition, they can act, whether favorably or not, on various mitochondrial processes. Some polyphenols have even been described to act on the signaling pathways (i.e. modulating intrinsic apoptosis (Mertens-Talcott et al., 2003; Zhang et al., 2013)) and trigger mitochondrial biogenesis (i.e. inducing sirtuins) (Baur et al., 2006; Chung et al., 2010; Davis et al., 2009; Sandoval-Acuna, Ferreira and Speisky, 2014a).

The notion that hesperidin behaves in a similar manner as quercetin is supported by several studies that show that quercetin supplementation also increases exercise performance and endurance capacity (Davis et al., 2010; MacRae and Mefferd, 2006). In particular, quercetin is also known to increase mitochondrial biogenesis and



exercise tolerance (Davis et al., 2009; Sandoval-Acuna et al., 2014b). While both hesperidin and quercetin exhibit mild uncoupling activity *in vitro* (Dorta et al., 2005), it is expected that, *in vivo*, hardly any uncoupling activity of flavonoids can be anticipated (van Dijk et al., 2000). This would increase proton leak and maximal respiratory capacity, supporting the hypothesis that hesperitin induces mitochondrial biogenesis.

Another point of interest is the effect of flavonoids on mitochondrial  $\text{Ca}^{2+}$  uniporter within the cell, increasing mitochondrial  $\text{Ca}^{2+}$  concentration.  $\text{Ca}^{2+}$  regulates several pathways, including one which stimulates eNOS, contributing to an increase of NO production, and stimulating  $\text{K}^{+}$  efflux (and  $\text{Ca}^{2+}$  influx), causing hyperpolarization of the cell membrane in endothelial cells (Garland et al., 2011). These results in relaxation of the smooth muscle in blood vessels and with this lowered blood pressure and increased blood flow (Reid, 2013; Umans and Levi, 1995; Wylie et al., 2013). Also, increased mitochondrial  $\text{Ca}^{2+}$  might stimulate oxidative metabolism by upregulation of pyruvate dehydrogenase and increase of mitochondrial membrane potential, allowing generation of high ATP concentrations lowering oxidative stress and muscle damage, ultimately increasing strength output (Bugger and Abel, 2010; Jouaville et al., 1999; Rutter and Rizzuto, 2000). The strong increase in power output in the present study (Table 2; Figure 2) is particularly interesting as it seems in contrast with typical hypotheses that mainly an improved blood flow, and therefore better oxygen delivery to the muscles, can improve exercise performance in athletes (Andersen and Saltin, 1985; Newman et al., 2002). Indeed, the observation that the  $\text{VO}_{2\text{max}}$  remains unchanged in combination with increased power output supports the potential role of the CF extract in the oxygen maintenance process in the muscle itself, rather than just increased oxygen delivery.

Furthermore, the fact that the positive results were obtained in a study in which well-trained athletes (average 9.6 hours per week) were included, further increases the relevance of the increase in power output. Indeed, it becomes exceedingly more difficult for trained individuals to gain further increase in strength and performance compared to untrained individuals (Ahtiainen et al., 2003; Peterson et al., 2005). An increase of 5% in overall power output is comparable with observed increases after creatine use (Buford et al., 2007). Interestingly, peak power output showed a trend to increase from baseline to 4 weeks after intervention in CF compared to placebo (Table 2). Although not significant ( $p = 0.112$ ) due to a large variation between participants, a calculated Cohen's  $d$  effect size of 2.70 shows a large magnitude of the treatment in the CF extract supplementation group on peak power output. An increase in peak power output through CF supplementation could be relevant for sports requiring immediate power generation.

To account for potential training adaptation, a familiarization test was conducted before the start of the study. But still, habituation with the test makes the athlete perform better along the study, simply because he is prepared better. Furthermore, daily exercise volume or nature (strength vs. endurance) was not recorded or standardized

by participants, although the nature of training has different effect on muscle alterations and metabolism (Baars, 2006). Also, several measurements that would have had added value to the study were not performed to lower the burden for the participant. For example,  $\text{VO}_{2\text{max}}$  testing could give more insight in aerobic fitness. Or, muscle biopsies would be of interest regarding mitochondrial number and function, for example to determine mitochondrial fractional area, an important quantitative indicator of oxidative capacity. Blood collection prior to and after exercise would have allowed for measurement of multiple blood markers for example related to oxidative stress, such as plasma malondialdehyde, reduced glutathione or cardiac troponin T, a marker for cardiac damage (Bloomer et al., 2005; Rifai et al., 1999). Both blood and muscle biopsies would have allowed a more mechanistic point of view, but were not included in this study due to their invasive character. Nevertheless, the study was placebo-controlled and the heart rate and  $\text{VO}_2$  did not change after 4 weeks, making it unlikely that the observed effects were a result of an overall increased fitness or extra effort specifically for the CF extract study group.

## Conclusions

Repeated intake of a specific CF extract may improve exercise performance in trained athletes. Therefore, this study is highly relevant for athletes to maximize their performance capacity. The improved performance might be due to improved respiratory efficiency in the mitochondria as a result of CF extract supplementation. However, further mechanistic research is needed to confirm this.

## Acknowledgements

The human intervention study was designed by JW and SP and executed by EO and JW. Data interpretation and manuscript preparation were performed by KW, EO, SP, and JL. The authors thank Ir. J. M. Schultink and Ir. A.B. Post (Team NL Innovation Center, Sportcentrum Papendal) for their technical assistance and Mr. T. Sam-sin (Wageningen University) for valuable discussions & suggestions. BioActor BV provided financial support to cover the costs of the study. KW and SP were employed by the sponsor company of the clinical trial at the time of study. The experiments comply with the current laws of the country in which they were performed. The authors have no conflict of interest to declare.

## References

- Ahtiainen, J.P., Pakarinen, A., Alen, M., Kraemer, W.J. and Hakkinen, K. (2003) Muscle hypertrophy, hormonal adaptations and strength development during strength training in strength-trained and untrained men. *European Journal of Applied Physiology* **89**(6), 555-563.
- Andersen, P. and Saltin, B. (1985) Maximal perfusion of skeletal muscle in man. *Journal of Physiology* **366**, 233-249.
- Baars, K. (2006) Training for Endurance and Strength: Lessons from Cell Signaling. *Medicine & Science in Sports & Exercise*, **38**(11), 1939-1944.
- Baur, J.A., Pearson, K. J., Price, N.L., Jamieson, H.A., Lerin, C., Kalra, A., Prabhu, V.V., Allard, J.S., Lopez-Lluch, G., Lewis, K., Pistell, P.J., Poosala, S., Becker, K.G., Boss, O., Gwinn, D., Wang, M., Ramaswamy, S., Fishbein, K.W., Spencer, R.G., Lakatta, E.G., Le Couteur, D., Shaw, R.J., Navas, P., Puigserver, P., Ingram, D.K., de Cabo, R. and Sinclair, D.A. (2006) Resveratrol improves health and survival of mice on a high-calorie diet. *Nature* **444**(7117), 337-342.

- Bescós, R., Sureda, A., Tur, J.A. and Pons, A. (2012) The effect of nitric-oxide-related supplements on human performance. *Sports Medicine* **2**, 99-117.
- Bloomer, R.J. (2010) Nitric oxide supplements for sports. *Strength & Conditioning Journal* **2**, 14-20.
- Bloomer, R.J., Goldfarb, A.H., Wideman, L., McKenzie, M.J. and Consitt, L.A. (2005) Effects of acute aerobic and anaerobic exercise on blood markers of oxidative stress. *Journal of Strength and Conditioning Research* **19**(2), 276-285.
- Bowtell, J.L., Summers, D.P., Dyer, A., Fox, P. and Mileva, K.N. (2011) Montmorency cherry juice reduces muscle damage caused by intensive strength exercise. *Medicine & Science in Sports & Exercise* **43**(8), 1544-1551.
- Buford, T.W., Kreider, R.B., Stout, J.R., Greenwood, M., Campbell, B., Spano, M., Ziegenfuss, T., Lopez, H., Landis, J. and Antonio, J. (2007) International Society of Sports Nutrition position stand: creatine supplementation and exercise. *Journal of the International Society of Sports Nutrition* **4**, 6.
- Bugger, H. and Abel, E.D. (2010). Mitochondria in the diabetic heart. *Cardiovascular Research* **88**(2), 229-240.
- Burke, L.M. and Read, R.S.D. (1993) Dietary Supplements in Sport. *Sports Medicine* **15**(1), 43-65.
- Chung, S., Yao, H., Caito, S., Hwang, J.-W., Arunachalam, G. and Rahman, I. (2010) Regulation of SIRT1 in cellular functions: Role of polyphenols. *Archives of Biochemistry and Biophysics*, **501**(1), 79-90.
- Cooke, M., John, P., Dzau, M. and Victor, J. (1997) Nitric oxide synthase: role in the genesis of vascular disease. *Annual Review of Medicine* **1**, 489-509.
- Davis, J.M., Carlstedt, C.J., Chen, S., Carmichael, M.D. and Murphy, E.A. (2010) The dietary flavonoid quercetin increases VO<sub>2</sub>(max) and endurance capacity. *International Journal of Sport Nutrition and Exercise Metabolism* **20**(1), 56-62.
- Davis, J.M., Murphy, E.A., Carmichael, M.D. and Davis, B. (2009) Quercetin increases brain and muscle mitochondrial biogenesis and exercise tolerance. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology* **296**(4), R1071-R1077.
- Dorta, D.J., Pigoso, A.A., Mingatto, F.E., Rodrigues, T., Prado, I.M., Helena, A.F., Uyemura, S.A., Santos, A.C. and Curti, C. (2005b) The interaction of flavonoids with mitochondria: effects on energetic processes. *Chemico-Biological Interactions* **152**(2-3), 67-78.
- Fradkin, A.J., Zazryn, T.R. and Smoliga, J.M. (2010) Effects of warming-up on physical performance: a systematic review with meta-analysis. *The Journal of Strength & Conditioning Research* **24**(1), 140-148.
- Froiland, K., Koszewski, W., Hingst, J. and Kopecky, L. (2004) Nutritional Supplement Use among College Athletes and Their Sources of Information. *International Journal of Sport Nutrition and Exercise Metabolism* **14**(1), 104-120.
- Garland, C.J., Hiley, C.R. and Dora, K.A. (2011) EDHF: spreading the influence of the endothelium. *British Journal of Pharmacology* **164**(3), 839-852.
- Gomez-Cabrera, M.-C., Domenech, E. and Viña, J. (2008) Moderate exercise is an antioxidant: Upregulation of antioxidant genes by training. *Free Radical Biology and Medicine* **44**(2), 126-131.
- Halliwell, B. (1991) Reactive oxygen species in living systems: Source, biochemistry, and role in human disease. *The American Journal of Medicine* **91**(3, Supplement 3), S14-S22.
- Jouaville, L.S., Pinton, P., Bastianutto, C., Rutter, G.A. and Rizzuto, R. (1999) Regulation of mitochondrial ATP synthesis by calcium: Evidence for a long-term metabolic priming. *Proceedings of the National Academy of Sciences of the United States of America* **96**(24), 13807-13812.
- MacRae, H.S. and Mefferd, K.M. (2006) Dietary antioxidant supplementation combined with quercetin improves cycling time trial performance. *International Journal of Sport Nutrition and Exercise Metabolism* **16**(4), 405-419.
- Malaguti, M., Angeloni, C. and Hrelia, S. (2013) Polyphenols in Exercise Performance and Prevention of Exercise-Induced Muscle Damage. *Oxidative Medicine and Cellular Longevity*, **2013**, 825928.
- Masella, R., Di Benedetto, R., Vari, R., Filesi, C. and Giovannini, C. (2005) Novel mechanisms of natural antioxidant compounds in biological systems: involvement of glutathione and glutathione-related enzymes. *The Journal of Nutritional Biochemistry* **16**(10), 577-586.
- McLeay, Y., Barnes, M.J., Mundel, T., Hurst, S.M., Hurst, R.D. and Stannard, S.R. (2012) Effect of New Zealand blueberry consumption on recovery from eccentric exercise-induced muscle damage. *Journal of the International Society of Sports Nutrition* **9**(1), 19.
- Mertens-Talcott, S. U., Talcott, S. T., & Percival, S. S. (2003). Low concentrations of quercetin and ellagic acid synergistically influence proliferation, cytotoxicity and apoptosis in MOLT-4 human leukemia cells. *Journal of Nutrition*, **133**(8), 2669-2674.
- Myburgh, K. H. (2014). Polyphenol supplementation: benefits for exercise performance or oxidative stress? *Sports Medicine, Suppl*, S57-70.
- Newman, J.M.B., Rattigan, S. and Clark, M.G. (2002) Nutritive blood flow improves interstitial glucose and lactate exchange in perfused rat hindlimb. *American Journal of Physiology - Heart and Circulatory Physiology* **283**(1), H186-H192.
- Nieman, D.C., Williams, A.S., Shanely, R.A., Jin, F., McAnulty, S.R., Triplett, N.T., Austin, M.D., and Henson, D.A. (2010) Quercetin's influence on exercise performance and muscle mitochondrial biogenesis. *Medicine & Science in Sports & Exercise* **42**(2), 338-345.
- Peternejl, T.-T. and Coombes, J.S. (2011) Antioxidant Supplementation during Exercise Training. *Sports Medicine* **41**(12), 1043-1069.
- Peterson, M.D., Rhea, M.R. Alvar, B.A. (2005) Applications of the dose-response for muscular strength development: a review of meta-analytic efficacy and reliability for designing training prescription. *Journal of Strength & Conditioning Research* **19**(4), 950-958.
- Powers, S.K., Nelson, W.B. and Hudson, M.B. (2011) Exercise-induced oxidative stress in humans: Cause and consequences. *Free Radical Biology and Medicine* **51**(5), 942-950.
- Reid, K. (2013) Performance Food: Promoting foods with a functional benefit in sports performance. *Nutrition Bulletin* **38**(4), 429-437.
- Rifai, N., Douglas, P.S., O'Toole, M., Rimm, E. and Ginsburg, G.S. (1999) Cardiac troponin T and I, electrocardiographic wall motion analyses, and ejection fractions in athletes participating in the Hawaii Ironman Triathlon. *The American Journal of Cardiology* **83**(7), 1085-1089.
- Rizza, S., Muniyappa, R., Iantorno, M., Kim, J.A., Chen, H., Pullikotil, P., Senese, N., Tesaro, M., Lauro, D., Cardillo, C. and Quon, M.J. (2011) Citrus polyphenol hesperidin stimulates production of nitric oxide in endothelial cells while improving endothelial function and reducing inflammatory markers in patients with metabolic syndrome. *Journal of Clinical Endocrinology & Metabolism* **96**(5), E782-E792.
- Rutter, G.A. and Rizzuto, R. (2000) Regulation of mitochondrial metabolism by ER Ca<sup>2+</sup> release: an intimate connection. *Trends in Biochemical Sciences* **25**(5), 215-221.
- Sachdev, S. and Davies, K.J. (2008) Production, detection, and adaptive responses to free radicals in exercise. *Free Radical Biology and Medicine* **44**(2), 215-223.
- Sandoval-Acuna, C., Ferreira, J. and Speisky, H. (2014a) Polyphenols and mitochondria: an update on their increasingly emerging ROS-scavenging independent actions. *Archives of Biochemistry and Biophysics* **559**, 75-90.
- Sandoval-Acuna, C., Ferreira, J. and Speisky, H. (2014b) Polyphenols and mitochondria: an update on their increasingly emerging ROS-scavenging independent actions. *Arch Biochem Biophys* **559**, 75-90.
- Santos, A.C., Uyemura, S.A., Lopes, J.L.C., Bazon, J.N., Mingatto, F.E. and Curti, C. (1998) Effect of Naturally Occurring Flavonoids on Lipid Peroxidation and Membrane Permeability Transition in Mitochondria. *Free Radical Biology and Medicine* **24**(9), 1455-1461.
- Trombold, J.R., Barnes, J.N., Critchley, L. and Coyle, E.F. (2010) Ellagitannin consumption improves strength recovery 2-3 d after eccentric exercise. *Medicine & Science in Sports & Exercise* **42**(3), 493-498.
- Trombold, J.R., Reinfeld, A.S., Casler, J.R. and Coyle, E.F. (2011) The effect of pomegranate juice supplementation on strength and soreness after eccentric exercise. *The Journal of Strength & Conditioning Research* **25**(7), 1782-1788.
- Tscholl, P., Alonso, J.M., Dollé, G., Junge, A. and Dvorak, J. (2010)

The Use of Drugs and Nutritional Supplements in Top-Level Track and Field Athletes. *The American Journal of Sports Medicine* **38**(1), 133-140.

- Umans, J.G. and Levi, R. (1995) Nitric oxide in the regulation of blood flow and arterial pressure. *Annual Review of Physiology* **57**, 771-790.
- van Dijk, C., Driessen, A.J. and Recourt, K. (2000) The uncoupling efficiency and affinity of flavonoids for vesicles. *Biochem Pharmacol* **60**(11), 1593-1600.
- Wylie, L.J., Mohr, M., Krstrup, P., Jackman, S. R., Ermidis, G., Kelly, J., Black, M.I., Bailey, S.J., Vanhatalo, A. and Jones, A.M. (2013) Dietary nitrate supplementation improves team sport-specific intense intermittent exercise performance. *European Journal of Applied Physiology* **7**, 1673-1684.
- Zhang, J.Y., Yi, T., Liu, J., Zhao, Z.Z. and Chen, H.B. (2013) Quercetin induces apoptosis via the mitochondrial pathway in KB and KBv200 cells. *Journal of Agricultural and Food Chemistry* **61**(9), 2188-2195.

### Key points

- Oxygen-processing capacity of mitochondria in the muscles plays a key role in exercise performance and recovery.
- In a double-blind, randomized, parallel study with 39 healthy, trained males the effect of 500 mg of a customized citrus flavonoid extract (CF) on exercise performance was assessed.
- CF intake significantly increased absolute power output with 5% and significantly decreased the oxygen consumption/power ratio, as  $VO_{2max}$  remained unchanged.
- The improved performance therefore may be due to improved respiratory efficiency as a result of CF extract supplementation, however, further mechanistic research is needed to confirm this.

## AUTHOR BIOGRAPHY

### Elvera OVERDEVEST

#### Employment

Researcher at Amsterdam University of Applied Sciences, Amsterdam, The Netherlands.

#### Degree

MSc

#### Research interest

The field of health promotion within the public health.

**E-mail:** e.j.overdevest@hva.nl

### Jeroen WOUTERS

#### Employment

Innovation Manager Sports and Nutrition at Sportcentrum Papendal, Arnhem, The Netherlands.

#### Degree

PhD

#### Research interest

Sports and nutrition.

**E-mail:** Jeroen.wouters@papendal.nl

### Kevin WOLFS

#### Employment

Clinical Research Associate at Novartis Pharma BV, Maastricht, The Netherlands.

#### Degree

MSc

#### Research interest

Oncology, nutritional and movement sciences.

**E-mail:** kevinwolfs@gmail.com

### Job van LEEUWEN

#### Employment

Research trainee at BioActor BV, Maastricht, The Netherlands

#### Degree

MSc

#### Research interest

Sports nutrition and exercise physiology.

**E-mail:** jjmvanleeuwen.bioactor@gmail.com

### Sam POSSEMIERS

#### Employment

R&D Director at BioActor BV, Maastricht, The Netherlands

#### Degree

PhD

#### Research interest

Bioactive potential of polyphenols and in gut health, gut barrier and immunity.

**E-mail:** sam.possemiers@bioactor.com

### ✉ Sam Possemiers

BioActor BV, Oxfordlaan 70, 6229 EV Maastricht, The Netherlands