

Research article

## Physiological Response Differences between Run and Cycle High Intensity Interval Training Program in Recreational Middle Age Female Runners

Milos Mallol<sup>1,2</sup>, Lynda Norton<sup>1</sup>, David J. Bentley<sup>3</sup>, Gaizka Mejuto<sup>4</sup>, Kevin Norton<sup>5</sup> and Javier Yanci<sup>2</sup>✉

<sup>1</sup> Exercise Science, College of Nursing and Health Sciences, Flinders University, South Bedford Park, South Australia; <sup>2</sup> Department of Physical Education and Sports, Faculty of Education and Sport, University of Basque Country (UPV-EHU), Portal de Lasarte Kalea, Vitoria-Gasteiz, Spain; <sup>3</sup> School of Environmental and Life Sciences Faculty of Science, University of Newcastle, Callaghan NSW, Australia; <sup>4</sup> Department of Body Expression Didactics, University of the Basque Country, Sarriena Auzoa. Leioa, Spain; <sup>5</sup> School of Health Sciences, City East Campus, University of South Australia, Adelaide, South Australia

### Abstract

The aim of this investigation was to compare the changes in endurance running performance and physiological variables after a four-week period of high intensity interval training (HIIT) in either running or cycling in female athletes. Fourteen recreational female runners (age =  $42 \pm 10$  yr, height =  $1.67 \pm 0.06$  m, body mass =  $61.6 \pm 10.4$  kg, body mass index (BMI) =  $22.2 \pm 3.4$  kg·m<sup>-2</sup>) were randomly allocated to one of two HIIT training groups: running (HIIT<sub>run</sub>) or cycling (HIIT<sub>bike</sub>). Each group performed two HIIT sessions per week for 4 weeks, which consisted of 6 x 2 min at 95% of maximal heart rate (HR<sub>max</sub>) and 4 x 1 min all out efforts. Maximal oxygen consumption (VO<sub>2max</sub>) in treadmill running increased significantly after the HIIT<sub>run</sub> ( $p < 0.01$ , ES = 0.6) but remained unchanged in HIIT<sub>bike</sub>. However, HIIT<sub>bike</sub> improved average velocity in a 10 km running time trial (TT<sub>run</sub>) ( $p < 0.05$ , ES = -0.4), whereas, no changes were found for the HIIT<sub>run</sub> group. Analysing the first and last HIIT sessions, for HIIT<sub>run</sub> only the average rate of perceived exertion (RPE<sub>av</sub>) increased significantly, whereas, performance variables such as average heart rate (HR<sub>av</sub>) and average pace (pace<sub>av</sub>) remained unchanged. HIIT<sub>bike</sub> enhanced significantly the average speed of HIIT sets (speed<sub>av</sub>) and the peak power output (PPO) of the session, as well as, the RPE<sub>av</sub> and delayed onset muscle soreness immediately after HIIT session (DOMS<sub>post</sub>) were increased significantly. A regime of HIIT in cycling may evoke increases in female recreational runners' power, which may be related with improvements in a 10 km TT<sub>run</sub> independent of changes in aerobic capacity. This may be advantageous in order to avoid overuse running related injuries.

**Key words:** gender, intermittent training, muscle damage, aerobic capacity, endurance.

### Introduction

High intensity interval training (HIIT) has been widely studied in athletic and non-athletic populations, demonstrating considerable positive benefits, such as, improved training time efficiency relative to increases in aerobic and anaerobic capacity or maintenance of endurance performance during low-volume training periods (Billat et al., 2001; García-Pinillos et al., 2017; Gunnarsson and Bangsbo, 2012; Lindsay et al., 1996). HIIT comes in many different forms and aims to improve distinct sport performances (Laursen and Jenkins, 2002). A number of authors

have investigated the optimal HIIT stimulus for improvements in endurance performance such as the intensity and duration of sessions (Etxebarria et al., 2014; Gunnarsson and Bangsbo, 2012; Laursen and Jenkins, 2002; Stepto et al., 1999). Despite the fact that HIIT appears to enhance a number of physiological variables, such as, maximal oxygen consumption (VO<sub>2max</sub>), peak power output (PPO), time to exhaustion at maximal velocity, first and second ventilatory threshold and vertical jumping performance (García-Pinillos et al., 2017; Helgerud et al., 2007; Laursen et al., 2002; Laursen and Jenkins, 2002; Mallol et al., 2018) the majority of research examining the effects of HIIT has been conducted with male athletes (Billat et al., 1999; Laursen et al., 2002; Lindsay et al., 1996) or with mixed male and female athletes sample (Farley et al., 2016; Koral et al., 2018; Mallol et al., 2016; Menz et al., 2015) while a group of exclusively female participants is less frequently studied. Some studies have reported less beneficial outcomes to HIIT training in females compared to males and hypothesized this may be due to a greater disposition to aerobic metabolism in females compared to males (Gibala et al., 2014; Gratas-Delamarche et al., 1994). Previous authors supported that gender might be a differentiating element in endurance performance, such as cross-country skiing, cycling and marathon running (Gibala et al., 2014; Gratas-Delamarche et al., 1994). Other studies have concluded that VO<sub>2max</sub>, maximal aerobic power, power at lactate threshold (LT), power at onset blood lactate accumulation (OBLA) and peak of speed levels showed significant differences between genders. Such that the male subjects obtained greater values (Hopker et al., 2010; Reaburn et al., 2011; Sandbakk et al., 2012). Additionally, anthropometric and morphological variables such as lipid accumulation may also be an influential factor (Reaburn et al., 2011).

The limited studies on female athletes have shown greater improvements in aerobic and anaerobic capacity after a HIIT program compared with continuous training in female soccer players (Rowan et al., 2012). The authors emphasised the time saving benefits obtained from HIIT sessions, helping to focus on teamwork and sport specific skills (Rowan et al., 2012). Kinnunen et al. (2017) found that a HIIT program applied in pre-season helped to enhance the maximal and explosive strength capacity, improving neuromuscular performance in female ice-hockey

players. Recently, Mallol et al. (2018) observed short supra-maximal sets of HIIT enhanced maximal  $\text{VO}_{2\text{max}}$  values and submaximal power [W] at the first and second ventilatory threshold capacity. However, no performance improvements were observed during a triathlon race simulation (Mallol et al., 2018). The context of the study was in relation to maintenance of fitness during periods of reduced training. There is minimal research examining the effects of HIIT in females. Therefore, further research is needed to achieve a greater understanding of this training method in females.

Previous researchers concluded that an intensified period of training in running results in a higher level of cumulative fatigue, greater muscle damage and potential injury in runners (Burt et al., 2012; Del Coso et al., 2013). An intensified training program, high volume and/or intensity, presents a demanding stimulus requiring careful planning and monitoring. Running can lead to higher levels of muscle damage and cumulative fatigue owing to eccentric muscle contractions which is evidenced by higher biochemical and perceptual markers of muscle damage and soreness such as creatine kinase (CK) (Burt et al., 2012; Cipryan, 2017; Keane et al., 2015; Quinn and Manley, 2012). In particular, muscle damage is typically observed in the periods after HIIT sessions. Elevated levels of CK, myoglobin and lactate dehydrogenase were induced by 15 s, 30 s and 60 s HIIT running protocols (Cipryan, 2017). Moreover, the muscular eccentric component of down-hill running training involved higher CK values and greater exercise induced inflammatory responses in female runners (Köhne et al., 2016). In female sports teams such as soccer, rugby and netball, female athletes presented an increased level of muscle damage after high intensity sprint training (Le Meur et al., 2011). Intensified HIIT in cycling, which is a more concentric based activity than running, may not result in the same cumulative level of soreness. Therefore, the current study focused on the comparing the residual effects of HIIT in cycling and running modalities. Additionally, Burt et al. (Burt et al., 2012) have presented data showing that different levels of exercise-induce muscle damage were evident following running and cycling exercises. However, as a general form of training HIIT in cycling might nonetheless induce performance changes similar to those shown following running HIIT programs (Burt et al., 2012; Millet et al., 2002) but this has not been extensively investigated. Several studies in well trained runners have observed positive effects on running performance when a part of their run training volume was replaced by cycle training sessions (Etxebarria et al., 2014; Tanaka, 1994; White et al., 2003). Overall, relatively few investigations have been conducted with female athletes in order to examine how an intensified running training program affects CK, as a biomarker of muscle damage. There is also limited research showing how exercise induced CK varies between running and cycling HIIT modes after a period of HIIT in running or cycling.

Therefore, the current investigation focused on examining the difference in physiological responses, performance outcomes and muscle damage, as acute effects,

occurring between run and cycle HIIT modes in female athletes.

## Methods

### Participants

A group of fourteen recreational middle-aged female athletes (age =  $42 \pm 10$  yr, height =  $1.67 \pm 0.06$  m, body mass =  $61.6 \pm 10.4$  kg, body mass index (BMI) =  $22.2 \pm 3.4$   $\text{kg}\cdot\text{m}^{-2}$ ) were recruited from a number of community clubs and institutions. Participants were randomly distributed in two groups: a running HIIT group (HIIT<sub>run</sub>, n = 7, age =  $41 \pm 7$  yr, height =  $1.64 \pm 0.07$  m, body mass =  $60.7 \pm 9.3$  kg, BMI =  $22.6 \pm 2.3$   $\text{kg}\cdot\text{m}^{-2}$ ) who completed two run HIIT sessions per week, and a cycling HIIT group (HIIT<sub>bike</sub>, n = 7, age =  $43 \pm 13$  yr; height =  $1.70 \pm 0.03$  m; body mass =  $62.5 \pm 12.1$  kg; BMI =  $21.8 \pm 4.5$   $\text{kg}\cdot\text{m}^{-2}$ ) who performed an identical HIIT session protocol on the cycle ergometer. The inclusion criteria were: participants were habitual and active runners,  $\geq$  two running sessions per week, and were able to run 10 km in  $< 70$  min. Participants were excluded if they had no running training in the previous one month, or had an injury that prevented them from participating in training or testing.

Participants were informed of the protocols and experimental procedures and signed a formal written consent. The study followed the guidelines established by the Declaration of Helsinki (2013) and was approved by the Human Research Ethics Committee (HREC code 334.16).

### Procedures

The study examined the physiological and performance benefits, as well as, the muscle damage generated by four weeks of HIIT using running or cycling in female runners. Participants attended Flinders University Exercise Physiology Laboratory twice weekly at the same time of day, to perform the HIIT sessions. To determine differences between the HIIT programs, participants were divided into two groups: HIIT<sub>run</sub> who performed supervised running HIIT sessions on an outdoor grass running track and HIIT<sub>bike</sub> who performed supervised cycling ergometer HIIT sessions in the laboratory. All participants performed identical testing procedures before and after the four-week intervention period. Each subject undertook the initial and final tests (laboratory testing and 10 km running time trial) at approximately the same time of day and were asked to follow a similar protocol for test preparation. The environmental conditions in the laboratory were maintained between 20–22°C and 55–65% humidity.

### Laboratory incremental running test

Each participant completed a 'fast' incremental exercise test to exhaustion using treadmill running in the week prior to and a week after the training intervention. The test procedure included (a) 10 minutes of their usual warm-up intensity on the treadmill (b) an incremental running test where the initial stage was set at  $8 \text{ km}\cdot\text{h}^{-1}$ . Each stage lasted one min and the speed was increased by  $1 \text{ km}\cdot\text{h}^{-1}$  until exhaustion (Noakes et al., 1990). Expired gases were ana-

lysed by TrueOne2400 (ParvoMedics, Utah, USA), to determine maximal oxygen consumption ( $VO_{2max}$ ) on a treadmill, the gas was analysed every 5 seconds but the 2 highest consecutive values over 30s was used. Before the warm up, blood lactate concentration was obtained from a fingertip sample and analysed using a portable lactate analyser (Lactate Pro, Arkray, KDK Corporation, Kyoto, Japan). Heart rate (HR), maximal speed achieved during the last stage completed ( $speed_{max}$ ) and rating of perceived exertion (RPE) were recorded in the final 15 s of every stage. HR was recorded using Polar RS400 series (Kempele, Finland) HR monitor until test completion. The Borg Scale (from 0 to 10) (Borg, 1982) was employed to monitor the RPE at the end of each stage. Participants were previously familiarised to the perceived exertion method and the 10-point Borg Scale.

### Running time trial (TT)

After a 60 min break from the incremental running test the participants completed a 10 km TT individually on a 400 m grass running track. Distance was previously measured and marked on the track by the researchers, participants were asked to complete 25 laps. Average and maximum speed ( $speed_{av}$  and  $speed_{max}$ ), and HR ( $HR_{av}$  and  $HR_{max}$ ) were recorded (Garmin Forerunner 910XT Olathe, Kansas, USA). RPE average ( $RPE_{av}$ ) was calculated from values obtained every 2 km throughout the trial. Instantaneous pace and HR were recorded every 2 km. Immediately after the test, lactate concentration was measured from a fingertip sample (Lactate Pro, Arkray, KDK Corporation, Kyoto, Japan).

### Training intervention

Participants randomly allocated to the HIIT groups ( $HIIT_{run}$  and  $HIIT_{bike}$ ) attended supervised training sessions twice a week. The session structure was similar for all HIIT training: a 10 min warm-up where participants determined their warm-up intensity from the maximal test, followed by 6 x 2 min at 95% of maximal heart rate ( $HR_{max}$ ) and 4 x 1 min all out (Bogdanis et al., 1996) followed by a 5 min cool-down. The recovery periods for the 1 and 2 min interval sets were 1 min 30 s and 2 min, respectively, with athletes continuing active recovery at a low intensity.  $HIIT_{run}$  performed the training outdoors around the same track as used for the 10 km TT. Individual training intensity at 95% of  $HR_{max}$  and maximal intensity for the  $HIIT_{run}$  group were determined for each participant based on their pre-intervention testing. The  $HIIT_{bike}$  sessions were performed using a cycle ergometer (Wattbike, Nottingham, UK). The  $HIIT_{bike}$  intensity were extrapolated from the treadmill test based on Millet et al. (Millet et al., 2009) review were researchers concluded that  $HR_{max}$  obtained from a maximal cycle ergometer test is about 5% (between 6-10 bpm) lower

than  $HR_{max}$  recorded in a maximal treadmill test. The intensity was also corroborated with RPE values (Basset and Boulay, 2000; Norton et al., 2010) which showed a similar pattern for HR during cycling interval training (Green et al., 2006). Participants could change the cycle resistance as they needed as long as they achieved the stipulated HR. In order to recognise their usual training load, they were asked to complete a 1-week training diary before the intervention to ensure that no significant differences existed between participants. During intervention weeks participants recorded their individual training sessions outside of the HIIT program. Every session was documented in a personal training diary which included the running training duration, distance and  $RPE_{av}$  allowing for the session RPE-min to be calculated (Table 1).

### Acute response to HIIT

The physiological responses to the first (1<sup>st</sup>) and last HIIT (8<sup>th</sup>) session in each mode ( $HIIT_{run}$  and  $HIIT_{bike}$ ) were measured. Blood lactate (Lactate Pro Analyser, Arkray, Japan) and CK concentrations (Reflotron Plus system, Rotkreuz, Switzerland) were measured before and immediately after the training session from fingertip blood samples. Participants came back to the laboratory 24 h after the HIIT sessions in order to measure  $CK_{24h}$  (Quinn and Manley, 2012). Delayed onset muscle soreness (DOMS) was recorded using a CR-10 scale (Lau et al., 2015) before, after and 24 h after session. During each  $HIIT_{run}$  session the average and peak HR, HR during recovery intervals, pace, distance covered and average/maximal RPE values were recorded (Borg, 1982). During the  $HIIT_{bike}$  average and maximal HR, recovery HR intervals and average and maximal RPE values (Borg, 1982) were recorded. Speed, power and cadence average, maximal power and cadence were also recorded for the  $HIIT_{bike}$  group.

### Statistical analyses

Results are presented as mean  $\pm$  SD. A t-test for independent samples was used to analyze the differences between  $HIIT_{run}$  and  $HIIT_{bike}$  at baseline (pre-test). The between-group ( $HIIT_{run}$  vs  $HIIT_{bike}$ ) comparison from pre-test to post-test or 1<sup>st</sup> and 8<sup>th</sup> HIIT sessions for data obtained in the laboratory tests and 10 km TT was calculated using a 2-way mixed ANOVA (group x time). In addition, a t-test for paired samples was used to analyze the differences between the pre-test and post-test independently for each group ( $HIIT_{run}$  or  $HIIT_{bike}$ ). Cohen's effect size (Cohen, 1988) was calculated to assess a practical significance between the pre-test and post-test in each group. Effect sizes (ES) of above 0.8, between 0.8 and 0.5, between 0.5 and 0.2, and lower than 0.2 were considered as large, moderate, small, and trivial, respectively (Cohen, 1988). Data analysis was performed using the Statistical Package for Social

**Table 1.** Quantification of training sessions, excluding HIIT sessions, for Run ( $HIIT_{run}$ ) and Bike ( $HIIT_{bike}$ ) groups.

	$HIIT_{run}$	$HIIT_{bike}$	p	Dif. (%)	ES
Duration (min)	49.97 $\pm$ 33.06	50.89 $\pm$ 23.99	0.84	1.84	0.0
Running distance (km)	6.59 $\pm$ 4.59	7.84 $\pm$ 4.22	0.12	18.99	0.3
$RPE_{av}$	4.31 $\pm$ 1.92	4.92 $\pm$ 1.90	0.06	14.28	0.3
Session RPE-min (AU)	227.53 $\pm$ 217.80	261.27 $\pm$ 166.36	0.27	14.83	0.1

Dif. (%) = mean differences in percentage, ES = effect size,  $RPE_{av}$  = average rate of perceived exertion, Session RPE-min = control method based on rate of perceived exertion per minute.

**Table 2. Results for Run (HIIT<sub>run</sub>) and Bike (HIIT<sub>bike</sub>) group in pre and post-intervention testing in maximal oxygen consumption (VO<sub>2max</sub>), maximum heart rate (HR<sub>max</sub>) and maximum speed (Speed<sub>max</sub>) obtained during maximal treadmill testing.**

	HIIT <sub>run</sub>					HIIT <sub>bike</sub>				
	Pre test	Post test	p	Dif. (%)	ES	Pre test	Post test	p	Dif. (%)	ES
VO <sub>2max</sub> (mL·kg <sup>-1</sup> ·min <sup>-1</sup> )	42.1 ± 4.9	45.2 ± 5.2**#	0.00	3.1 ± 1.6	0.6	41.0 ± 5.3	41.8 ± 6.5	0.26	0.8 ± 1.5	0.1
HR <sub>max</sub> (bpm)	187 ± 8	181 ± 11	0.09	-5.9 ± 7.7	-0.5	174 ± 9	173 ± 10	0.16	-1.6 ± 2.6	-0.2
Speed <sub>max</sub> (km·h <sup>-1</sup> )	14.1 ± 1.4	14.6 ± 1.6	0.11	0.4 ± 0.6	0.3	13.8 ± 1.5	14.3 ± 1.9	0.20	0.5 ± 0.8	0.3

Dif. (%) = mean differences in percentage, ES = effect size. \*\* p < 0.01 significant differences with pre test. # p < 0.05, two way mixed ANOVA analysis (group x time) statistical differences.

Sciences (SPSS Inc, version 24.0 for Windows, Chicago, IL, USA). The statistical significance was set at p < 0.05. Despite the fact that in some cases, a variable showed a p value > 0.05, whereas, ES was greater than 0.5 (moderate), was considered practical difference.

## Results

No significant differences were found pre-intervention between HIIT<sub>run</sub> and HIIT<sub>bike</sub> for VO<sub>2max</sub>, HR<sub>max</sub> and Speed<sub>max</sub> in the incremental treadmill test or any variable obtained during the 10 km TT. After 4-weeks of the HIIT intervention, HIIT<sub>run</sub> improved VO<sub>2max</sub> significantly (p < 0.01, ES = 0.6, moderate), and decreased HR<sub>max</sub> (p = 0.09, ES = -0.5, moderate) whereas the Speed<sub>max</sub> values were maintained (p = 0.11, ES = 0.3, small). In the HIIT<sub>bike</sub>, no changes were observed after 4-weeks for VO<sub>2max</sub>, HR<sub>max</sub> and Speed<sub>max</sub> (p = 0.16 to 0.26, ES = -0.2 to 0.3, trivial to small) (Table 2). According to the two-way mixed ANOVA analysis (group x time), only VO<sub>2max</sub> showed a statistically significant difference. The HIIT<sub>run</sub> group enhanced the VO<sub>2max</sub> result in the post-test (p = 0.01) while the HIIT<sub>bike</sub> group showed no change.

After 4-weeks of the intervention, neither HIIT<sub>run</sub> nor HIIT<sub>bike</sub> changed either the time to complete the 10 km TT (p = 0.06 to 0.84), ES = 0.1 to -0.2, trivial to small), or average heart rate (HR<sub>av</sub>) (p = 0.14 to 0.58, ES = -0.2 to -0.3, small) (Table 3). HIIT<sub>run</sub> resulted in a high lactate concentration at the end of the test (Lactate<sub>post</sub>) (p = 0.18 to 0.05, ES = 3.6, very large). There was a significant increase in the average and maximum rating of perceived exertion (RPE<sub>av</sub>) (p = 0.04, ES = 1.7, large) and RPE<sub>max</sub> (p = 0.04, ES = 1.6, large), whereas, HR<sub>av</sub> did not change significantly (p > 0.05, ES = -0.2, trivial) (Table 3). In the HIIT<sub>bike</sub> group the average pace decreased significantly (p = 0.02, ES = -0.3 small) and the maximal rate of perceived exertion (RPE<sub>max</sub>) increased during the 10 km TT post-test (p = 0.04, ES = 0.9, large) (Table 3). There were no group x time differences.

**Table 3. Results for Run (HIIT<sub>run</sub>) and Bike (HIIT<sub>bike</sub>) group in pre and post- intervention tests for, lactate values, average pace (Pace<sub>av</sub>), Time, heart rate average (HR<sub>av</sub>), maximum heart rate (HR<sub>max</sub>), average rating of perceived exertion (RPE<sub>av</sub>) and maximum rate of perceived exertion (RPE<sub>max</sub>) obtained in the 10 km time trial.**

	HIIT <sub>run</sub>					HIIT <sub>bike</sub>				
	Pre test	Post test	p	Dif. (%)	ES	Pre test	Post test	p	Dif. (%)	ES
Lactate (mmol/l)	3.3 ± 0.8	6.6 ± 0.9	0.18	3.3 ± 0.6	3.6	5.9 ± 2.5	7.8 ± 4.4	0.23	1.8 ± 3.6	0.4
Pace <sub>av</sub> (min/km)	5.36 ± 0.8	5.36 ± 0.7	0.91	0.0 ± .5	0.0	5.48 ± 1.3	5.30 ± 1.0*	0.02	-0.4 ± 0.3	-0.4
Time (min)	57.4 ± 7.3	57.8 ± 6.0	0.84	-0.4 ± 4.5	0.1	59.7 ± 11.9	57.5 ± 10.8	0.06	-2.1 ± 2.5	-0.02
HR <sub>av</sub> (bpm)	172 ± 9	170 ± 8	0.58	-1.4 ± 6.5	-0.2	169 ± 10	166 ± 12	0.14	-3.6 ± 5.5	-0.3
RPE <sub>av</sub> (AU)	5 ± 2	7 ± 1*	0.05	1.5 ± 1.6	1.7	6 ± 2	6 ± 2	0.41	0.4 ± 1.3	0.3
RPE <sub>max</sub> (AU)	7 ± 2	8 ± 1*	0.04	1.3 ± 1.3	1.6	6 ± 1	8 ± 2*	0.05	1.6 ± 1.7	0.9

Dif. (%) = mean differences in percentage, ES = effect size. \*p < 0.05 significant differences with pre test

Table 3 and Figure 1 show the results for the 1<sup>st</sup> and 8<sup>th</sup> HIIT sessions. HIIT<sub>run</sub> demonstrated a significant increase for RPE<sub>av</sub> (p = 0.03, ES = 0.8, moderate) and an increase in lactate concentration values immediately after the session (Lactate<sub>post</sub>) (p = 0.14, ES = 0.6, moderate) (Table 4) and creatine-kinase (CK) concentration before session (CK<sub>pre</sub>) (p = 0.15, ES = 1.0, large) (Figure 1). No significant differences were observed for the remaining variables. The HIIT<sub>bike</sub> group showed a significant improvement in Speed<sub>av</sub>, P<sub>av</sub>, and P<sub>max</sub> (p = 0.01 to 0.03, ES = 0.6 to 0.7, moderate), while increases in RPE<sub>av</sub> and delayed onset muscle soreness immediately after training (DOMS<sub>post</sub>) were also observed (p = 0.02, ES = 1.4 to 2.2, large) (Table 4). The maximal heart rate obtained during HIIT sets and recovering sets (HR<sub>max-work</sub> and HR<sub>maxrecovery</sub>) and Lactate<sub>post</sub> demonstrated practical increases although these were not significant (p = 0.16 to 0.34, ES = 0.6 to 2.0, moderate to large). The average heart rate recorded during recovery intervals (HR<sub>av recovery</sub>) and the concentration of CK 24-hour after the HIIT session (CK<sub>24after</sub>) decreased practically (p = 0.46 to 0.49, ES = -0.6 to -1.0, moderate to large) (Figure 1). According to the two-way mixed ANOVA analysis (group x time), only DOMS<sub>post</sub> showed statistical significance (p = 0.017).

## Discussion

Previous studies have analysed the effects of HIIT in individual exercise modes on a single mode of activity (Cook et al., 2010; Laursen et al., 2002; Sijie et al., 2012). However, the effects of this training method in running or cycling on performance in a single mode such as running in female athletes has not been studied. In the current investigation, only the HIIT<sub>run</sub> group evoked significant improvement in VO<sub>2max</sub> during a maximal treadmill test. Neither HIIT<sub>run</sub> nor HIIT<sub>bike</sub> significantly enhanced the time to complete 10 km TT, despite a significant decrease in average pace for the HIIT<sub>bike</sub> participants. Running HIIT generated a significantly greater level of muscular fatigue in non-

**Table 4.** Results in 1<sup>st</sup> and 8<sup>th</sup> HIIT session for Run HIIT (HIIT<sub>run</sub>) and for Bike HIIT (HIIT<sub>bike</sub>) group.

	1 <sup>st</sup> HIIT	8 <sup>th</sup> HIIT 8	p	Dif. (%)	ES	
HIIT <sub>run</sub>	HR <sub>av-work</sub> (bpm)	167 ± 5	166 ± 10	0.91	-0.3 ± 6.4	-0.0
	HR <sub>max-work</sub> (bpm)	182 ± 6	183 ± 10	0.69	1.0 ± 5.7	0.1
	HR <sub>av-recovery</sub> (bpm)	148 ± 9	147 ± 17	0.91	-0.4 ± 9.6	-0.0
	HR <sub>max-recovery</sub> (bpm)	181 ± 7	181 ± 10	1.00	0.0 ± 4.2	0.0
	Pace <sub>av</sub> (min·km <sup>-1</sup> )	4.11 ± 0.5	3.59 ± 0.4	0.27	-0.1 ± 0.3	-0.4
	Distance <sub>av</sub> (m)	386.2 ± 22.5	389.8 ± 30.6	0.54	3.6 ± 11.9	0.1
	RPE <sub>av</sub> (AU)	7 ± 1	8 ± 2*	0.03	1.1 ± 1.1	0.8
	RPE <sub>max</sub> (AU)	10 ± 1	10 ± 1	1.00	0.0 ± 1.0	0.0
	Lactate <sub>pre</sub> (U·L <sup>-1</sup> )	1.7 ± 0.4	1.6 ± 0.7	0.74	-0.1 ± 0.6	-0.1
	Lactate <sub>post</sub> (U·L <sup>-1</sup> )	6.3 ± 2.2	8.0 ± 3.2	0.14	1.8 ± 2.8	0.6
	DOMS <sub>pre</sub> (AU)	0.6 ± 1.5	1.0 ± 2.2	0.20	0.4 ± 0.8	0.2
	DOMS <sub>post</sub> (AU)	2.0 ± 1.6	2.4 ± 1.9	0.43	0.4 ± 1.1	0.2
	DOMS <sub>24After</sub> (AU)	1.0 ± 2.4	1.2 ± 2.4	0.36	0.2 ± 0.4	0.1
HIIT <sub>bike</sub>	HR <sub>av-work</sub> (bpm)	153 ± 6	152 ± 12	0.77	-1.3 ± 11.0	-0.1
	HR <sub>max-work</sub> (bpm)	155 ± 23	164 ± 13	0.16	9.2 ± 12.0	0.7
	HR <sub>av-recovery</sub> (bpm)	145 ± 15	139 ± 11	0.49	-6.2 ± 20.3	-0.6
	HR <sub>max-recovery</sub> (bpm)	156 ± 18	168 ± 6	0.29	12.2 ± 22.3	2.0
	Speed <sub>av</sub> (km·h <sup>-1</sup> )	33.9 ± 4.5	36.2 ± 3.0*	0.02	2.3 ± 1.8	0.7
	P <sub>av</sub> (W)	166 ± 49	198 ± 44**	0.01	32.4 ± 20.0	0.7
	P <sub>max</sub> (W)	240 ± 77	280 ± 63*	0.03	40.0 ± 36.8	0.6
	Cadence <sub>av</sub> (rpm)	93 ± 12	93 ± 6	0.84	0.7 ± 9.0	0.1
	Cadence <sub>max</sub> (rpm)	103 ± 13	104 ± 8	0.86	0.7 ± 9.9	0.1
	RPE <sub>av</sub> (AU)	7 ± 1	8 ± 1*	0.02	1.7 ± 1.5	2.2
	RPE <sub>max</sub> (AU)	9 ± 1	10 ± 1	0.09	0.7 ± 1.0	1.5
	Lactate <sub>pre</sub> (mmol·l <sup>-1</sup> )	1.8 ± 0.6	1.5 ± 0.6	0.39	-0.3 ± 0.8	-0.5
	Lactate <sub>post</sub> (mmol·l <sup>-1</sup> )	10.3 ± 3.8	11.6 ± 2.3	0.34	1.3 ± 3.4	0.6
	DOMS <sub>pre</sub> (AU)	0.3 ± 0.8	2.0 ± 2.2	0.12	1.7 ± 2.5	0.8
	DOMS <sub>post</sub> (AU)	1.9 ± 1.1	6.1 ± 3.0*#	0.02	4.3 ± 3.4	1.4
DOMS <sub>24After</sub> (AU)	2.3 ± 2.2	1.6 ± 1.4	0.14	-0.7 ± 1.1	-0.5	


Variables: Work interval average Heart rate (HR<sub>av-work</sub>), Work interval maximum Heart rate (HR<sub>max-work</sub>), recovery interval average Heart rate (HR<sub>av-recovery</sub>), recovery interval maximum Heart rate (HR<sub>max-recovery</sub>), average pace (Pace<sub>av</sub>), work average distance (Distance<sub>av</sub>), average speed (Speed<sub>av</sub>), Power average (P<sub>av</sub>), Maximum Power obtained (P<sub>max</sub>), Average cadence (Cadence<sub>av</sub>), maximum cadence (Cadence<sub>max</sub>), average work interval rate of perceived exertion (RPE<sub>av</sub>) and work interval maximum rate of perceived exertion (RPE<sub>max</sub>), Lactate concentration pre test (Lactate<sub>pre</sub>), lactate concentration post test (Lactate<sub>post</sub>), Delayed Onset Muscle soreness pre test (DOMS<sub>pre</sub>), Delayed Onset Muscle soreness post test (DOMS<sub>post</sub>), Delayed Onset Muscle soreness 24 hour after test (DOMS<sub>24After</sub>). Dif. (%) = mean differences in percentage, ES = effect size. \*p < 0.05, \*\* p < 0.01 significant differences with pre test. # p < 0.05, two way mixed ANOVA analysis (group x HIIT session) statistical differences.

competitive female runners compared to cycling HIIT.

As noted above, VO<sub>2max</sub> improved significantly in the HIIT<sub>run</sub> group while the remaining maximal test variables in each training group were not significantly modified. This situation may be due in part to an excessive training amount accumulated by runners who may not have been accustomed to structured HIIT. Previous investigations observed improvements in VO<sub>2max</sub> and maximal test variables such as HR and speed or power, however the performance level changed from the current investigation. Rowan et al. (2012) observed an enhancement of 4.73% in VO<sub>2max</sub> after 5-week of 5 x 30 s with 4.5 min recovery in female soccer players, however, similar results were obtained from the control group who performed 40 min continuous running at 80% of VO<sub>2max</sub>. In another study, Gunnarsson and Bangsbo (2012) showed that 7-weeks of short interval HIIT improved VO<sub>2max</sub> in moderately trained male and female runners. Mallol et al. (2018) concluded that after 4-weeks of a cycling HIIT program in moderately trained triathletes, participants improved 6.7% in VO<sub>2max</sub> and 15% in peak power. Finally, Lesmes et al. (1978) after 8-weeks of two types of supra maximal interval training (short duration at 170% of velocity at VO<sub>2max</sub> (vVO<sub>2max</sub>) and long duration at 130% at vVO<sub>2max</sub>) concluded that the frequency of training, interval distances and intensities were independ-

ent of changes in aerobic power and submaximal HR in females, whereas, interval training intensity was essential to improve these variables in males rather than frequency and interval distance. In our study, HIIT<sub>run</sub> achieved greater changes than HIIT<sub>bike</sub> and this may be due to the specificity of the activity. For future investigations, it would be relevant to consider the effects of cycling HIIT during a maximal test using a cycle ergometer. Additionally, comparing different HIIT interventions of different intensity and work to rest characteristics.

During the 10 km TT the HIIT<sub>run</sub> performance remained unchanged but RPE<sub>av</sub> increased significantly indicating a greater perception of effort. At the same time, HR<sub>max</sub> decreased after 4-week of HIIT<sub>run</sub> program. Perceived exertion may have increased during post-intervention running 10 km TT because of individual fatigue accumulated after the HIIT program possibility due to insufficient recovery after the period of intensified training. Controversially, other researchers have observed improvements in running performances after a HIIT program. Gunnarsson and Bangsbo (2012) obtained an improvement of 6% and 4% in 1500 m and 5 km running tests, respectively. The protocol employed included 7-weeks of interval running working at intensities of 90% of HR<sub>max</sub> with a 54% training volume reduction in moderately trained females.



**Figure 1. Comparison between A) HIIT<sub>run</sub> and B) HIIT<sub>bike</sub> for CK pre, immediately post and 24-hour after HIIT session.** Variables: Creatine Kinase values pre session (pre); Creatine Kinase values immediately post session (post); CK values 24 hour after the session (24 after); First HIIT session of the program (1st HIIT); last session of the program (8th HIIT).

In addition Paavolainen et al. (1999) showed an improvement in 5 km run time in well-conditioned athletes with no changes in  $VO_{2max}$ , similar to that observed in the current study results for HIIT<sub>bike</sub> participants. Furthermore, Bangsbo et al. (2009) observed a decrease of 1 min in 10 km performance (from 37 min to 36 min) after a 6-9-week training period with intervals near maximal speed with a 30% reduced total training volume. However, Iaia et al. (2009) determined that after 4-week of 8-12 x 30 s maximal speed intervals and a 64% reduction in total training volume, but there were no improvements in 10 km TT run in these endurance trained participants. The studies mentioned above enhanced running performance after HIIT interventions, however, a number of these studies replaced a part of the participants' usual training volume with HIIT sessions. In the current research, the intensities employed during HIIT intervals were maximal or close to maximal which may have generated an excessive training stress, and therefore, excess residual fatigue to the amateur female runners.

By contrast, the HIIT<sub>bike</sub> participants significantly increased performance ( $Pace_{av}$ , min/km) during the 10 km TT run with an increased  $RPE_{max}$ . Similarly, Mikesell and Dudley (1984) noted a decrease of 81 s on 10 km distance in well-trained runners after an intensive aerobic program, combining 40 min run "all out" 3 days a week with 5 x 5 min at  $VO_{2max}$  intervals with 5 min jogging on the treadmill as recovery. A number of investigations conducted with cyclists concluded that cycling HIIT significantly enhances cycling performance in a range of testing protocols and competitive simulations (Laursen et al., 2002; Lindsay et al., 1996; Stepto et al., 1999). The use of HIIT using identical activity modes to that of competition helped to improve the sports performance. Despite the fact that the HIIT<sub>bike</sub> group did not show significant differences for time to complete 10 km TT, the  $Pace_{av}$  manifested a significant decrease, hence, performance during the 10 km TT post-test was enhanced. Whilst the mechanisms accounting for this performance improvement are unknown, it is possible that a gain in lower limb muscular power because of the

cycle HIIT sessions led to this improvement. It is also possible that the exercise mode of running presented a higher physiological demand due to a greater muscle recruitment, contraction type and accumulative fatigue which was associated with the different performance adaptations between modes (Le Meur et al., 2011). Nevertheless, the present study results suggest that the HIIT<sub>run</sub> program did not result in an improvement in 10 km performance considering that a greater lactate<sub>post</sub> concentration and RPE<sub>max</sub> were observed post-test despite a similar HR<sub>av</sub> and similar time to complete the time trial.

This study is novel in that biochemical and perceptive markers associated with muscle soreness were measured in the acute stages following the first (1<sup>st</sup>) and last (8<sup>th</sup>) HIIT sessions. Following the HIIT<sub>run</sub> intervention RPE<sub>av</sub>, lactate<sub>post</sub>, and specifically, CK<sub>pre</sub> were elevated, whereas, distance completed and Pace<sub>av</sub> remained the same from the 1<sup>st</sup> HIIT compared with the 8<sup>th</sup> HIIT session. This potentially indicates a greater level of muscle damage and lack of assimilation to the HIIT running program. By contrast, HIIT<sub>bike</sub> participants improved Speed<sub>av</sub> and Power<sub>av</sub> from the 1<sup>st</sup> to 8<sup>th</sup> HIIT session. This enhancement was associated with more positive responses for RPE<sub>av</sub>, RPE<sub>max</sub>, DOMS<sub>post</sub>, lactate<sub>post</sub> and HR<sub>max</sub>, although HR<sub>av</sub> decreased during recovery, suggesting that HIIT<sub>bike</sub> athletes were able to perform more effectively in the session. At the same time both groups showed greater CK<sub>pre</sub> during the 8<sup>th</sup> HIIT. However, the difference between sessions were smaller, once again this may have occurred due to fatigue accumulation and lack of assimilation of the high intensity training. Furthermore, CK<sub>24h</sub> values after the 8<sup>th</sup> HIIT session decreased compared with the 1<sup>st</sup> HIIT in the HIIT<sub>bike</sub> group, which may show a physiological adaptation such that HIIT<sub>bike</sub> runners were able to achieve a superior level of recovery from muscular fatigue.

Previous studies have focused on markers of muscle damage such as, CK concentration and DOMS after a high intensity training program in runners and cyclist while performing running and cycling testing in isolation (Nieman et al., 2014). In these studies, it was shown that muscle damage was related to activity mode. Nieman et al. (2014) analysed differences between running and cycling performances on runners and cyclists after 3 consecutive days of an ‘overreaching’ training program, concluding that the eccentric contractions intrinsic to running, resulted in 133% greater CK concentration and 87% greater DOMS in runners immediately after 3 days of running training compared with cyclists performing similar training on a cycle ergometer. Additionally, CK concentrations remained more elevated in runners than in cyclists 1, 14 and 38 h after the training program. Similarly, DOMS presented a comparable pattern to CK at 1 and 14 h and by 38 h post intervention; the values were similar in both groups. Likewise, Bruunsgaard et al. (1997) and Proske and Allen (2005) concluded that eccentric activities generated higher levels of muscle damage (CK, DOMS and myoglobin) compared with isometric and concentric contractions. In our study, CK<sub>pre</sub> before the first and last HIIT were greater in the cycling group than the running group. However, the difference between sessions for CK<sub>post</sub> were similar in both groups. For its part, CK<sub>24h</sub> results showed that HIIT<sub>bike</sub>

might be able to recover faster than HIIT<sub>run</sub> after a HIIT program.

One of the current research limitations relates to sample size. Longitudinal studies, which require the completion of exhaustive and multiple assessments during an intervention, are onerous on participants, particularly taking into account work and family commitments. This can make it difficult to recruit large numbers of participants and contributed to some of the drop out in the current study. Another limitation relates to the timing of the 10 km TT on the same day of, and following, the maximal incremental test, the completion of multiple, exhaustive tests may have detrimentally affected the performance outcomes for subsequent tests on the same day. This is somewhat ameliorated in that all participants undertook the same testing process both before and after the intervention.

## Conclusion

After 4-weeks of a HIIT program, only HIIT<sub>run</sub> participants improved VO<sub>2max</sub> whereas no improvements were observed for 10 km TT run performance potentially due to fatigue accumulation generated by the HIIT training itself. Although no significant group by time differences were observed, the HIIT<sub>bike</sub> participants demonstrated improved 10 km TT run performance (Pace<sub>av</sub>) indicative of positive cross training transfer. This occurred without changes in VO<sub>2max</sub> during maximal incremental running tests. Both HIIT modes evoked some muscle damage although HIIT<sub>bike</sub> seems to have achieved faster muscular recovery 24 h after HIIT session completion. Therefore, it appears that a HIIT<sub>run</sub> program in recreational female athletes produces excessive stress, fatigue and muscle damage which may have resulted in inadequate stimulus for enhancement of running performance, whereas, HIIT<sub>bike</sub> may be a beneficial training modality that can be used to improve running performance.

## Acknowledgements

The experiments comply with the current laws of the country in which they were performed. The authors have no conflict of interest to declare.

## References

- Bangsbo, J., Gunnarsson, T.P., Wendell, J., Nybo, L. and Thomassen, M. (2009) Reduced volume and increased training intensity elevate muscle Na<sup>+</sup>-K<sup>+</sup> pump  $\alpha$ 2-subunit expression as well as short- and long-term work capacity in humans. *Journal of Applied Physiology* **107**, 1771-1780.
- Basset, F.A. and Boulay, M.R. (2000) Specificity of treadmill and cycle ergometer tests in triathletes, runners and cyclists. *European Journal of Applied Physiology* **81**, 214-221.
- Billat, V., Slawinski, J., Bocquet, V., Chassaing, P., Demarle, A. and Koralsztejn, J. (2001) Very Short (15 s-15 s) Interval-Training Around the Critical Velocity Allows Middle-Aged Runners to Maintain VO<sub>2</sub> max for 14 minutes. *International Journal of Sports Medicine* **22**, 201-208.
- Billat, V.L., Flechet, B., Petit, B., Muriaux, G. and Koralsztejn, J.-p. (1999) Interval training at VO<sub>2max</sub>: effects on aerobic performance and overtraining markers. *Medicine and Science in Sports and Exercise* **31**, 156-163.
- Bogdanis, G.C., Nevill, M.E., Boobis, L.H. and Lakomy, H. (1996) Contribution of phosphocreatine and aerobic metabolism to energy supply during repeated sprint exercise. *Journal of Applied Physiology* **80**, 876-884.
- Borg, G.A. (1982) Psychophysical bases of perceived exertion. *Medicine and Science in Sports and Exercise* **14**, 377-381.

- Bruunsgaard, H., Galbo, H., Halkjaer-Kristensen, J., Johansen, T., MacLean, D. and Pedersen, B. (1997) Exercise-induced increase in serum interleukin-6 in humans is related to muscle damage. *The Journal of Physiology* **499**, 833-841.
- Burt, D., Lamb, K., Nicholas, C. and Twist, C. (2012) Effects of muscle-damaging exercise on physiological, metabolic, and perceptual responses during two modes of endurance exercise. *Journal of Exercise Science & Fitness* **10**, 70-77.
- Cipryan, L. (2017) IL-6, antioxidant capacity and muscle damage markers following high-intensity interval training protocols. *Journal of Human Kinetics* **56**, 139-148.
- Cohen, J. (1988) *Statistical power analysis for the behavioral sciences*. Hillsdale, NJ: Lawrence Earlbaum Associates **2**.
- Cook, K., Cathcart, A.J., Scott, R.A. and Easton, C. (2010) High Intensity Interval Training increases aerobic and anaerobic capacity in collegiate female soccer players: 2643. *Medicine & Science in Sports & Exercise* **42**, 698-699.
- Del Coso, J., Fernández, D., Abián-Vicen, J., Salinero, J.J., González-Millán, C., Areces, F., Ruiz, D., Gallo, C., Calleja-González, J. and Pérez-González, B. (2013) Running pace decrease during a marathon is positively related to blood markers of muscle damage. *PLoS one* **8**, e57602.
- Etxebarria, N., Anson, J.M., Pyne, D.B. and Ferguson, R.A. (2014) High-intensity cycle interval training improves cycling and running performance in triathletes. *European Journal of Sport Science* **14**, 521-529.
- Farley, L.O.R., Secomb, L.J., Parsonage, R.J., Lundgren, E.L., Abbiss, R.C. and Sheppard, M.J. (2016) Five Weeks of Sprint and High-Intensity Interval Training Improves Paddling Performance in Adolescent Surfers. *Journal of Strength and Conditioning Research* **30**, 2446-2452.
- García-Pinillos, F., Cámara-Pérez, J.C., Soto-Hermoso, V.M. and Latorre-Román, P.Á. (2017) A High Intensity Interval Training (HIIT)-based running plan improves athletic performance by improving muscle power. *The Journal of Strength & Conditioning Research* **31**, 146-153.
- Gibala, M.J., Gillen, J.B. and Percival, M.E. (2014) Physiological and health-related adaptations to low-volume interval training: influences of nutrition and sex. *Sports Medicine* **44**, 127-137.
- Gratas-Delamarche, A., Le Cam, R., Delamarche, P., Monnier, M. and Koubi, H. (1994) Lactate and catecholamine responses in male and female sprinters during a Wingate test. *European Journal of Applied Physiology and Occupational Physiology* **68**, 362-366.
- Green, J.M., McLester, J.R., Crews, T.R., Wickwire, P.J., Pritchett, R.C. and Lomax, R.G. (2006) RPE association with lactate and heart rate during high-intensity interval cycling. *Medicine and Science in Sports and Exercise* **38**, 167-172.
- Gunnarsson, T.P. and Bangsbo, J. (2012) The 10-20-30 training concept improves performance and health profile in moderately trained runners. *Journal of Applied Physiology* **113**, 16-24.
- Helgerud, J., Hoydal, K., Wang, E., Karlsen, T., Berg, P., Bjerkaas, M., Simonsen, T., Helgesen, C., Hjørt, N. and Bach, R. (2007) Aerobic High-Intensity Intervals Improve VO<sub>2</sub>max More Than Moderate Training. *Medicine and Science in Sports and Exercise* **39**, 665.
- Hopker, J., Jobson, S., Carter, H. and Passfield, L. (2010) Cycling efficiency in trained male and female competitive cyclists. *Journal of Sports Science & Medicine* **9**, 332.
- Iaia, F.M., Hellsten, Y., Nielsen, J.J., Fernström, M., Sahlin, K. and Bangsbo, J. (2009) Four weeks of speed endurance training reduces energy expenditure during exercise and maintains muscle oxidative capacity despite a reduction in training volume. *Journal of Applied Physiology* **106**, 73-80.
- Keane, K.M., Salicki, R., Goodall, S., Thomas, K. and Howatson, G. (2015) Muscle damage response in female collegiate athletes after repeated sprint activity. *The Journal of Strength & Conditioning Research* **29**, 2802-2807.
- Kinnunen, J.-v., Piitulainen, H. and Piirainen, J.M. (2017) Neuromuscular adaptations to short-term high-intensity interval training in female ice hockey players. *Journal of Strength and Conditioning Research*.
- Köhne, J.L., Ormsbee, M.J. and McKune, A.J. (2016) The effects of a multi-ingredient supplement on markers of muscle damage and inflammation following downhill running in females. *Journal of the International Society of Sports Nutrition* **13**, 44.
- Koral, J., Oranchuk, D.J., Herrera, R. and Millet, G.Y. (2018) Six Sessions of Sprint Interval Training Improves Running Performance in Trained Athletes. *Journal of Strength and Conditioning Research* **32**, 617-623.
- Lau, W.Y., Blazevich, A.J., Newton, M.J., Wu, S.S.X. and Nosaka, K. (2015) Assessment of muscle pain induced by elbow-flexor eccentric exercise. *Journal of Athletic Training* **50**, 1140-1148.
- Laursen, P.B., Blanchard, M.A. and Jenkins, D.G. (2002) Acute high-intensity interval training improves Tvent and peak power output in highly trained males. *Canadian Journal of Applied Physiology* **27**, 336-348.
- Laursen, P.B. and Jenkins, D.G. (2002) The scientific basis for high-intensity interval training. *Sports Medicine* **32**, 53-73.
- Le Meur, Y., Hausswirth, C., Natta, F., Bignet, F. and Vidal, P. (2011) A multidisciplinary approach to overreaching detection in endurance trained athletes. *Journal of Applied Physiology* **114**, 411-420.
- Lesmes, G.R., Fox, E.L., Stevens, C. and Otto, R. (1978) Metabolic responses of females to high intensity interval training of different frequencies. *Medicine and Science in Sports* **10**, 229-232.
- Lindsay, F.H., Hawley, J.A., Myburgh, K.H., Schomer, H.H., Noakes, T.D. and Dennis, S.C. (1996) Improved athletic performance in highly trained cyclists after interval training. *Medicine and Science in Sports and Exercise* **28**, 1427-1434.
- Mallol, M., Bentley, D.J., Norton, L., Norton, K., Mejuto, G. and Yanci, J. (2018) Comparison of Reduced Volume-High Intensity Interval Training Compared to High Volume Training on Endurance Performance in Triathletes. *International Journal of Sports Physiology and Performance* 1-26.
- Mallol, M., Mejuto, G. and Bentley, D. (2016) Effects of 4 weeks high-intensity training on running and cycling performance in well-trained triathletes. *Sports and Exercise Medicine Open Journal* **2**, 55-61.
- Menz, V., Strobl, J., Faulhaber, M., Gatterer, H. and Burtcher, M. (2015) Effect of 3-week high-intensity interval training on VO<sub>2</sub>max, total haemoglobin mass, plasma and blood volume in well-trained athletes. *European Journal of Applied Physiology* **115**, 2349-2356.
- Mikesell, K.A. and Dudley, G.A. (1984) Influence of intense endurance training on aerobic power of competitive distance runners. *Medicine and Science in Sports and Exercise* **16**, 371-375.
- Millet, G.P., Candau, R., Barbier, B., Busso, T., Rouillon, J.-D. and Chatard, J.-C. (2002) Modelling the transfers of training effects on performance in elite triathletes. *International Journal of Sports Medicine* **23**, 55-63.
- Millet, G.P., Vleck, V.E. and Bentley, D.J. (2009) Physiological differences between cycling and running. *Sports Medicine* **39**, 179-206.
- Mujika, I. and Padilla, S. (2003) Scientific bases for precompetition tapering strategies. *Medicine & Science in Sports & Exercise* **35**, 1182-1187.
- Nieman, D.C., Luo, B., Dréau, D., Henson, D.A., Shanelly, R.A., Dew, D. and Meaney, M.P. (2014) Immune and inflammation responses to a 3-day period of intensified running versus cycling. *Brain, Behavior, and Immunity* **39**, 180-185.
- Noakes, T.D., Myburgh, K.H. and Schall, R. (1990) Peak treadmill running velocity during the V O<sub>2</sub> max test predicts running performance. *Journal of Sports Sciences* **8**, 35-45.
- Norton, K., Norton, L. and Sadgrove, D. (2010) Position statement on physical activity and exercise intensity terminology. *Journal of Science and Medicine in Sport* **13**, 496-502.
- Paavola, L., Hakkinen, K., Hamalainen, I., Nummela, A. and Rusko, H. (1999) Explosive-strength training improves 5-km running time by improving running economy and muscle power. *Journal of Applied Physiology* **86**, 1527-1533.
- Proske, U. and Allen, T.J. (2005) Damage to skeletal muscle from eccentric exercise. *Exercise and Sport Sciences Reviews* **33**, 98-104.
- Quinn, T.J. and Manley, M.J. (2012) The impact of a long training run on muscle damage and running economy in runners training for a marathon. *Journal of Exercise Science & Fitness* **10**, 101-106.
- Reaburn, P.R., Dascombe, B.J. and Janse de Jonge, X. (2011) Body composition and gender differences in performance. In: *Nutritional Assessment of Athletes*. Second Edition. Eds:



- Driskell JA, Wolinsky I, Eds CRC Press, Boca Raton, FL 121-147.
- Rowan, A.E., Kueffner, T.E. and Stavrianeas, S. (2012) Short duration high-intensity interval training improves aerobic conditioning of female college soccer players. *International Journal of Exercise Science* **5**, 6.
- Sandbakk, Ø., Ettema, G., Leirdal, S. and Holmberg, H.-C. (2012) Gender differences in the physiological responses and kinematic behaviour of elite sprint cross-country skiers. *European Journal of Applied Physiology* **112**, 1087-1094.
- Sijie, T., Hainai, Y., Fengying, Y. and Jianxiong, W. (2012) High intensity interval exercise training in overweight young women. *The Journal of Sports Medicine and Physical Fitness* **52**, 255-262.
- Stepito, N.K., Hawley, J.A., Dennis, S.C. and Hopkins, W.G. (1999) Effects of different interval-training programs on cycling time-trial performance. *Medicine and Science in Sports and Exercise* **31**, 736-741.
- Tanaka, H. (1994) Effects of Cross-Training: Transfer of Training Effects on  $\dot{V}O_{2\max}$  between Cycling, Running and Swimming. *Sports Medicine* **18**, 330-339.
- White, L.J., Dressendorfer, R.H., Muller, S.M. and Ferguson, M.A. (2003) Effectiveness of cycle cross-training between competitive seasons in female distance runners. *The Journal of Strength & Conditioning Research* **17**, 319-323.

### Key points

- This article observes the effects of a HIIT program in two different modes, i.e. cycling and running, in a hardly investigated population, recreational middle-age female runners.
- The analysis of physiological variables in a HIIT session, maximal test and 10 km time trial run performance.

### ✉ Javier Yanci

Department of Physical Education and Sports, Faculty of Education and Sport, University of Basque Country (UPV-EHU), Portal de Lasarte Kalea 71, 01007 Vitoria-Gasteiz, Spain

### AUTHOR BIOGRAPHY



#### Milos MALLO

##### Employment

PhD candidate at Flinders University (South Australia) and the University of the Basque Country/ UPV-EHU (Spain)

##### Degree

MSc

##### Research interests

Physiological and performance adaptations in endurance athletes.

**E-mail:** mall0078@flinders.edu.au

#### Lynda NORTON

##### Employment

Health and Exercise Science, College of Nursing and Health Sciences, Flinders University, South Bedford Park, South Australia

##### Degree

MSc

##### Research interests

Cardiopulmonary Exercise testing and training in clinical and sedentary populations.

**E-mail:** lynda.norton@flinders.edu.au

#### David J. BENTLEY

##### Employment

The program convenor, Master of Exercise Physiology and Rehabilitation at the Discipline of Exercise and Sport Science, University of Newcastle.

##### Degree

PhD

##### Research interests

Multidisciplinary areas in exercise and medical sciences.

**E-mail:** david.bentley@newcastle.edu.au

#### Gaizka MEJUTO

##### Employment

Lectureship and research at Department of Body Expression Didactics, University of the Basque Country, Spain

##### Degree

PhD

##### Research interests

Physiological adaptation of the human body to different environments for the best performance, athlete testing and monitoring.

**E-mail:** gaizka.mejuto@ehu.eus

#### Kevin NORTON

##### Employment

Professor of Exercise Science. He holds an academic position as a researcher and teacher at the University of South Australia.

##### Degree

PhD

##### Research interests

High-performance sport, evolution of athletes and the physiological limits of human capabilities.

**E-mail:** Kevin.Norton@unisa.edu.au



#### Javier YANCI

##### Employment

Senior Lecturer at University of the Basque Country (UPV/EHU).

##### Degree

PhD

##### Research interests

Sport performance analysis, high-performance sport and sport training methods.

**E-mail:** javier.yanci@ehu.eus