

Research article

Moderate and Severe Injuries at Five International Olympic-Style Wrestling Tournaments during 2016-2019

Szabolcs Molnár^{1,2,3} ✉, Zolt Hunya⁴, Krisztián Gáspár⁵, Imre Szerb^{6,7}, Noé Szabó⁸, Károly Mensch⁹, Éva Körösi¹⁰, Katalin Bacskai¹¹, Ákos Kálmán Sántha^{2,12}, Eszter Anna Janka⁵ and Babak Shadgan^{3,13}

¹ Department of Traumatology, Medical Centre Hungarian Defense Forces, Budapest, Hungary; ² University of Physical Education, Budapest, Hungary; ³ Medical, Prevention & Anti-Doping Commission of The United World Wrestling, Corsier-sur-Vevey, Switzerland; ⁴ Department of Orthopedics, University of Debrecen, Debrecen, Hungary; ⁵ Department of Dermatology, University of Debrecen, Debrecen, Hungary; ⁶ Department of Traumatology, Semmelweis University, Budapest, Hungary; ⁷ Department of Orthopaedics of Uzsoki Hospital, Budapest, Hungary; ⁸ Semmelweis University Medicine and Health Sciences, Budapest, Hungary; ⁹ Department of Oral Diagnostics, Semmelweis University, Faculty of Dentistry, Budapest, Hungary; ¹⁰ Dietitian and WADA coordinator of the Hungarian Wrestling Federation, Budapest, Hungary; ¹¹ National Institute for Sports Medicine, Budapest, Hungary; ¹² Professional Medical Services and Assistance Ltd., Budapest, Hungary; ¹³ Department of Orthopaedics, The University of British Columbia, Vancouver, Canada

Abstract

As a contact sport, wrestling may result in injuries. Based on the severity, they are classified as mild, moderate, severe and critical. All injuries occurring at international competitions are documented in a cloud-based surveillance system. The purpose of this study was to analyze the incidence and characteristics of moderate and severe (including critical) wrestling injuries that occurred during five international Olympic-style wrestling competitions in 2016-2019. Three Wrestling World Championships and two European Wrestling tournaments were organized by the Hungarian Wrestling Federation in 2016-2019. A total of 2483 wrestlers in three Olympic wrestling styles have competed in 3007 matches. Data from all injuries were recorded and analyzed to define rates, locations, types and severity, and to compare with previous reports. A total of 53 wrestlers sustained 55 injuries, which is equivalent to an overall injury incidence rate of 9.1‰ (9.1/1000 athletic exposures). Greco-Roman and Women Wrestling had the same injury incidence rate, while Freestyle had a lower one (9.5‰ versus 8.5‰). The injury proportion by regions and anatomic locations were on head and face 29.1%, spine and trunk 16.4 % and the upper-and-lower extremity injuries equally 27.3%. The most common types of injuries included ligament lesions, joint injuries, skin lacerations, and contusions. Five wrestlers (0.8‰) sustained strangulation or concussion. Wrestling injury rates during United World Wrestling competitions are not high, but when happen they can be serious. Despite relatively low incidence rate of injuries, there is a need for continuous education for medical teams, referees and coaches to avoid wrestling injuries.

Key words: Wrestling; injuries; head-and-trunk; extremities; strangulation; incidence

Introduction

Olympic wrestling is a high-demand contact sport, which occasionally results in injury (Halloran, 2008; Shadgan et al., 2010). Prevention and reduction of injuries are considered as main priorities by the United World Wrestling (UWW), the international governing body of Olympic styles wrestling. One of the main structures of this international sports federation is the Medical, Prevention

and Anti-Doping Commission (UWW-MC). Continuous upgrading and monitoring of the UWW Medical Regulations and direct supervision of the medical and doping control coverages of wrestling competitions are essential duties of the UWW-MC (UnitedWorldWrestling, 2019a). A Practical Medical Guide for Wrestling Competitions is also prepared to help those physicians or health care providers who are neither familiar with the wrestling rules nor with common wrestling injuries (Molnár et al., 2014). To understand and control wrestling injuries, all injuries and medical conditions during UWW competitions are recorded and analyzed by UWW-MC through a well-established secured cloud-based surveillance system, so-called “Athena”. This information helps to monitor and consider proper actions to prevent injuries and improve the safety and the healthcare of wrestlers (Channon et al., 2020).

Understanding the mechanisms of wrestling injuries is essential to prevent and reduce the risk of injuries; however, it is not always easy in wrestling due to the complexity of movements in this sport (Molnár et al., 2020; Myers et al., 2010). Besides frequent contacts and collisions during wrestling, inadequate preparation or loss of concentration can result in injuries too (Maffulli et al., 2010; Molnár et al., 2020). The dietary regimen may also contribute to physical injuries in wrestlers (Barley et al., 2019; Jlid et al., 2013; Khodaei et al., 2015; Kim and Park, 2021; Lingor and Olson, 2010). Though the bouts and training sessions take part in soft mats, tested and permitted by UWW and there is an increasing number of other precautions and improvements to defend wrestlers as well, the development of injuries are inevitable (Shadgan et al., 2010; Shadgan et al., 2021; Shadgan et al., 2017; Tomin and Kmetty, 2021).

The definition of injury in UWW competitions includes “any musculoskeletal or soft tissue complaint newly incurred during the competitions that required medical attention regardless of the consequences with respect to absence from sport” (Shadgan et al., 2010; Shadgan et al., 2017). The severity of wrestling injuries is

categorized by the UWW-MC as mild (treated on the mat), moderate (additional medical care is required after the competition at the venue), severe and critical (competition is terminated by the referee or the venue medical personnel and the injured wrestler should receive emergency care at an equipped medical centre and be immediately transferred to a nearby hospital). Any severe injury that places life in jeopardy is classified as Critical injury (UnitedWorldWrestling, 2019b).

The purpose of this study was to investigate the incidence and the characteristics of contact (direct and indirect) moderate, severe and critical injuries during wrestling competitions by direct observation in order to improve the Medical Service by the local Organizers and by the UWW.

Methods

Moderate, severe and critical injuries occurred during five UWW Wrestling Championships organized by the Hungarian Wrestling Federation from 2016 to 2019 were studied. These competitions included the 2016 and the 2018 Senior World Championships, the 2019 U23 (wrestlers between 18-23 years old) World Championships, the 2017 U23 European Championships and the 2018 U15 (wrestlers between 13-15 years old) European Championships. All competitions included three Olympic wrestling styles, Greco-Roman (GR), Freestyle (FS) and Women Wrestling (WW).

Theoretically one wrestler could compete in many different Championships thus from a statistical point of view we considered every entry individually in each competition.

A medical team consisted of seven health care professionals from the listed authors was in charge of medical coverage and injury data collection by direct observation during all five events. The documentation and the orientational diagnosis were recorded on-site, based on the injury pattern followed by physical examination.

All direct and indirect contact injuries were documented in UWW Injury Report Forms and uploaded into the secured cloud-based UWW injury surveillance system, Athena (UnitedWorldWrestling, 2019b). As there are many ways to quantify athletic exposure, we chose

wrestling specific adaptation of International Olympic Committee consensus statement about exposure measures (Bahr et al., 2020).

The collected data were used to calculate the rates of injuries and proportions in different styles and age groups, and to quantify the location, type and severity of injuries. The type of treatment and the continuation of competitions post injury were also studied. The data collected during these competitions was shared strictly for scientific research with the authors of this study and always kept confidential.

Statistical analysis

Chi-squared or Fisher's exact test were used to assess categorical variables. The influencing factors of the severity of the injuries (moderate vs. severe and critical) were determined by binomial logistic regression. The odds ratio (OR) was counted with corresponding 95% upper and lower confidence intervals (CI). Statistical significance was established at the 5.0% significance level. Statistical analysis was performed by SPSS 25 (SPSS package for Windows, Release 25; SPSS Inc., Chicago, Illinois, USA).

Ethical approval

This study was planned, conducted and realized on behalf of the Medical, Prevention and Anti-Doping Commission of the United World Wrestling.

Results

General injury occurrence

A total of 2483 wrestlers participated in 3007 wrestling matches, which accounts for 6014 athlete-exposures (AEs) as one match equals two athletes' exposures. A total of 55 moderate, severe and critical injuries in 53 wrestlers were documented, indicating an injury incidence rate of 9.1 injuries per 1000 AEs (Table 1).

Injuries according to wrestling styles and age groups

Although the proportion of injuries in different styles was different (from 27.3% to 38.2%), the injury incidence rate were very close, ranging from 8.5 to 9.5 injuries per 1000 AEs). However the proportion of injuries by age groups (from 12.7% to 58.2%) and the distribution of injury

Table 1. Occurrence of moderate and severe (including critical) injuries during World and Continental Championships organized by the Hungarian Wrestling Federation between 2016-2019.

Competition	athletes			Number of matches			Injuries (injured wrestler)			
	GR*	FS†	WW‡	GR	FS	WW	GR	FS	WW	
Senior World Championship (non-Olympic weight categories)	56	50	36	65	59	44	3	0	0	
U23 European Championship	147	116	91	172	137	107	3	3 (2)	4	
U15 European Championships	237	205	174	277	241	201	1	2	4	
Senior World Championships	296	266	225	343	306	261	6	4	3	
U23 Senior World Championships	217	219	148	255	363	176	8	10 (9)	4	
	953	856	674	1112	1106	789	21	19 (17)	15	
Sum		2483			3007			55 (53)		
Injury incidence rate / AEs§				6014 AEs				9.1‰ = 9.1/1000 AEs		

*Greco-Roman Wrestling, †Freestyle Wrestling, ‡Women Wrestling, §athletes' exposures

incidence rate (from 5‰ among U15, 13‰ in U23 and 7.5‰ in Seniors) demonstrated a more significant ranging (Table 2).

Injured body parts by regions and by anatomic location

The most common regions of injuries were the head and face (29.1% among the injured body parts with 2.6‰ injury incidence rate), followed by upper and lower extremities (27.3% and 27.3%, respectively – 2.5‰ per AEs). Spine and trunk were injured in 16.4% of the cases which equals to 1.5‰ injury incidence rate. Regarding the anatomic regions, the injuries of the extremities were the most common (54.5%), followed by the head, face, spine and trunk injuries (45.5%).

The most injured body part by anatomic location was the knee (21.8% of injury proportion and 2‰ per AEs), followed by the head (14.5% vs 1.3‰), the face (14.5% vs 1.3‰) and then the ribs (12.7% vs 1.2‰). Other injuries occurred in the shoulder, elbow, hand, fingers, ankle and lumbar spine. All injuries based on their regions and anatomical locations are listed in Table 3.

Type of injury

There were equal numbers of ligament lesions (n= 6 sprain & n= 6 rupture) and joint injuries (n= 3 dislocation, n= 3 subluxation & n= 6 sprain). Other injuries included skin laceration (n= 11), contusion (n= 7 ribs, n= 1 clavicle, n=1 forearm & n= 1 metacarpal), strangulation (n= 3) and

concussion (n=2). Muscle injuries, bone fracture and nerve injury were less common. Table 4 depicts the details of injury types including the injury incidence rate per AEs.

Injury severity

56.4% of all injuries were classified as severe (including one critical injury where a temporary paraplegia occurred), and 43.6% were moderate. Table 5 displays combined information of severity and location of injuries in different styles and age groups.

The influencing factors of injury severity (moderate vs. severe & critical) were analyzed by binomial logistic regression. Different variables including age group, wrestling style, weight category and injury location were included in the model. The analysis showed that there was a greater odd of severe injuries on the extremities (4.61 [1.26; 16.92]) (Table 6).

Discussion

Injury is a barrier to sports participation and development (Shadgan et al., 2017). Although wrestling injuries are comprehensively studied by investigators, the definition of injuries and the population of the studies are extremely heterogeneous (Hewett et al., 2005).

It is very difficult to compare injuries if the data search focused on different patterns, such as incidents that officially halted a match (Kersey and Rowan, 1983), or

Table 2. Injuries according to styles and age groups.

Style	Number of matches	Number of injuries	Proportion of injuries	Injury incidence rate / AEs§ (6014)
GR*	1112	21	38.2%	9.5‰
FS†	1106	19	34.4%	8.5‰
WW‡	789	15	27.3%	9.5‰
Age group	Number of matches	Number of injuries	Proportion of injuries	Injury incidence rate / AEs§ (6014)
Senior	1078	16	29.1%	7.5‰
U23	1210	32	58.2%	13‰
U15	719	7	12.7%	5‰

§athletes' exposures.

Table 3. Injured body parts - Number of injuries by regions and by anatomic locations.

Regions Anatomic locations	Number of injuries	Injury proportion by body parts	Injury incidence rate / AEs§ (6014)
Head & Face	16	29.1%	2.6‰
Head'	8	14.5%	1.3‰
Face''	8	14.5%	1.3‰
Spine & Trunk	9	16.4 %	1.5‰
Ribs	7	12.7%	1.2‰
Lumbar Spine	2	3.6%	0.3‰
Upper Extremity	15	27.3%	2.5‰
Shoulder joint	5	9.1%	0.8‰
Elbow	5	9.1%	0.8‰
Hand and finger	3	5.4%	0.5‰
Clavicle	1	1.8%	0.2‰
Forearm	1	1.8%	0.2‰
Lower Extremity	15	27.3%	2.5‰
Knee'''	12	21.8%	2‰
Ankle	2	3.6%	0.3‰
Groin	1	1.8%	0.2‰

§athletes' exposures. 'Head includes: 3 strangulations, 2 concussions and 3 skin lacerations. ''Face includes: forehead 3, ear 1, eyebrow 4, (all were skin lacerations, rest of the face were unharmed). '''Knee includes 1 knee cup (patellar injury).

Table 4. Type of injuries.

Type of Injury	Number	Injury proportion	Injury incidence rate / AEs§ (6014)
Ligament lesion (6 sprain & 6 rupture)	12	21.8%	2‰
Joint (3 dislocation, 3 subluxation & 6 sprain)	12	21.8%	2‰
Skin laceration	11	20%	1.8‰
Contusion (7 ribs, 1 clavicle, 1 forearm & 1 metacarpal)	10	18.2%	1.6‰
Strangulation (3) and concussion (2)	5	9%	0.8‰
Muscle	3	5.4%	0.5‰
Traumatic bone fracture	1	1.8%	0.2‰
Nerve injury	1	1.8%	0.2‰

§athletes' exposures

Table 5. Severity and Location of injury proportion by Style and by age groups.

Indicators	# Inj	# Inj %	Style - #Inj				Age groups - #Inj				
			GR*	FS†	WW‡	p-value	Senior	U23	U15	p-value	
Severity of injury	moderate	24	43.6%	11	8	5	0.516	6	15	3	0.924
	severe & critical	31	56.4%	10	11	10		10	17	4	
Location of injury	head-and-trunk	25	45.5%	11	13	1	0.001**	7	16	2	0.590
	upper-and-lower extremities	30	54.5%	10	6	14		9	16	5	

#|Number of injuries, *Greco-Roman Wrestling, †Freestyle Wrestling, ‡Women Wrestling. ** p<0.05

Table 6. Logistic regression.

Severity of injury (reference: moderate)	Variables	OR¶ [95% CI††]	p-value	
Severe & critical	Age group	U15/Senior	0.55 [0.06; 4.71]	0.584
		U23/Senior	0.59 [0.15; 2.26]	0.436
	Wrestling style	FS†/WW‡	0.58 [0.11; 3.07]	0.519
		GR*/WW	1.36 [0.23; 8.09]	0.739
	Weight category (kg)		1.00 [0.97; 1.04]	0.783
	Injury location	extremities/head and trunk	4.61 [1.26; 16.92]	0.021**

**p < 0.05. ¶|odds ratio. ††95% CI: 95% confidence intervals, *Greco-Roman Wrestling, †Freestyle Wrestling, ‡Women Wrestling

reached the athletic training room (Strauss and Lanese, 1982), or limitations of function to an extent that the athlete sought treatment by an athletic trainer or physician (Lorish et al., 1992; Pasque and Hewett, 2000) or restricted participation of at least one day beyond the initial injury (Kordi et al., 2012).

Another key issue is the location where the documentation is done, which can be recorded during the training, in the competition or at the hospital (Hewett et al., 2005; Otero et al., 2017; Yard et al., 2008). The time frame of data collection is also an important issue, whether it was assembled in one competition, or in a school-year, or life-long in a wrestling club or during a preparation period of a national team (Kordi et al., 2012; Otero et al., 2017; Park et al., 2019; Pasque and Hewett, 2000; Yamaner et al., 2012; Yard and Comstock, 2008). The accuracy of recorded injury data also varies in different studies. Of all the competition data are likely to be the most accurate because they are directly observed and recorded by medical teams (Thomas and Zamanpour, 2018). There are many ways to quantify athletic exposure and no single measure will suit all surveillance settings and research questions. The choice of exposure measures is heavily influenced by sport-specific and contextual factors, as well as which types of health problems are of interest. We adopted the International Olympic Committee consensus statement about exposure measures specifically to wrestling (Bahr et al., 2020).

We examined the direct and indirect contact injuries that occurred during these five international competitions documented by direct observation (Bahr et al., 2020). The

wrestling injury rates were 9.1/1000 AEs findings. Kroshus et al. (2018) analyzed 10 academic years where they found a similar injury incidence rate among college wrestlers, but a lesser rate in high school (9.28 versus 2.38/1000 AEs). Thomas and Zamanpour (2018) reported a higher injury incidence rate of 16.3/1000AE for competition studies and 69.5/1000AE for databases. Shadgan et al. (2010; 2017) studied all wrestling injuries that lead to discontinuation of the match in three Olympic Games. The moderate and severe cases from his data accounted for 6 to 12,5 injury incidence rates per 1000 AE from Beijing to Rio Olympic Games.

Regarding the injuries suffered in different styles Shadgan et al. (2017) concluded that during the Rio Olympic competitions the distribution of all wrestling injuries was 40.9% GR, 36.4% FS and 22.7% WW style respectively (Shadgan et al., 2017). Our data shows a similar injury proportion. (Estwanik et al., 1978; Thomas and Zamanpour, 2018; Yard and Comstock, 2008) all reported that GR style had lower injury rates compared to FS; however, the subject populations of those studies were different. Regarding the age groups Strauss and Lanese (1982) studied four wrestling tournaments in grade school, high school and college-age levels and found a 3.8% injury rate among grade schoolboys, who were injured less frequently than high school or college wrestlers (12%). These data also correlate to our findings.

In our study, the injuries of the upper-and-lower extremities were the most common, followed by the head-and-trunk injuries. Hewett and his colleagues (2005) reviewed 10 studies of junior wrestling injuries where the

body region incurring the greatest percentage of injuries was the head/spine/trunk (range of 24.5–48%, average 36.7%), followed by the upper (range of 9.3–42%, average 27.7%) and lower extremities (range of 7.5–45.1%, average 28.75%). Thomas and Zamanpour (Thomas and Zamanpour, 2018) collected data from eight studies. Their results weighed almost the same proportion as we found: 31% head and neck injuries, 25.7% upper extremities, 24.4% lower extremities and 15.4% trunk and spine injuries.

The knee was the far most injured body part by anatomic location in our research as it was in other studies (Jarrett et al., 1998; Kroshus et al., 2018). We found eight head and eight face injuries (21.9% of all injuries). (Kersey and Rowan, 1983) reported that 36.4% of injuries involved the head-face-neck region while Kroshus et al. (2018) reported 21.3% of head and face injuries (Kroshus et al., 2018). The number of rib injuries accounted for 12.7% in our study. Hewett et al. (2005) found rib and chest injuries ranging from 4.1% to 16.1% in different studies, however blunt trauma is difficult to diagnose and treat (Dzsinich et al., 2015). We found an equal number of five injuries in the shoulder and elbow (9.1% of injuries). Different authors reported higher rates of shoulder injuries (3.5 to 24%) and lower rates of elbow injuries (1 to 10.5%) (Agel et al., 2007; Hewett et al., 2005; Kroshus et al., 2018; Pasque and Hewett, 2000; Shadgan et al., 2017). Hand and finger injuries accounted for 5.45% of all injuries in our study, almost the same as Kroshus found (6%) (Kroshus et al., 2018). Ankle and lumbar spine injuries (2 cases each) were accounted for 3.6% in our study, while ankle had a much higher proportion rate in two different studies 7.5% (Agel et al., 2007) and 6.9% (Kroshus et al., 2018)). Lower back injuries varied from 1.2 to 18.6% (Agel et al., 2007; Hewett et al., 2005; Strauss and Lanese, 1982) in three different reviews. In our study, we recorded one injury (1.8%) for each of the clavicle, the forearm and the groin. Hewett et al. (2005) and Agel et al. (2007) ranged pelvis and hip injuries between 1.1–3.2% in their studies. Interestingly, we did not observe any dental injury in our competitions while dental and orofacial trauma is not rare in wrestling (Hewett et al., 2005; Kvittem et al., 1998).

In terms of the type of injuries there were almost equal numbers of ligament lesions, joint injuries, and skin laceration and contusion. Shadgan et al. (2017) reported skin laceration as the most frequent injuries (41%) followed by ligament sprains (13.6%), joint dislocations (13.6%) and contusions (9%) in wrestlers during Rio Olympic competitions. Thomas and Zamanpour (2018) have also reported sprain and strain to be the most frequent injury type with a 37.6% proportion rate which is higher than our findings, followed by laceration, abrasion and contusion which accounted for 23.4% altogether.

We recorded 3 cases of strangulation (5.4%) and 2 cases of concussion (3.6%). Shadgan et al. (2017) reported one case of strangulation injury (4.5%) during the Rio Olympic competitions (Shadgan et al., 2017). Kroshus et al. (2018), and Thomas and Zamanpour (2018) reported a higher rate of concussion (15.9% and 26%, respectively). Prevention of concussion that is due an immediate unintentional action can only be prevented by well-prepared wrestlers, while strangulation injury can be

avoided by cautious refereeing (Hewett et al., 2005; Yeo et al., 2020).

Muscle strain and traumatic bone fracture accounted for lesser incidence (5.4% -1.8% respectively) in our cohort while Agel et al. (2007) and Kroshus et al. (2018) reported higher rates of muscle-tendon strain for up to 13.2% -13.3%, respectively. Thomas and Zamanpour (2018) reviewed three articles (Agel et al., 2007; Pasque and Hewett, 2000; Yard and Comstock, 2008) about bone fractures during wrestling competitions and found a 5.8% injury rate.

In our study we observed a wrestler who suffered a strangulation injury followed by temporary paraplegia (nerve injury 1.8%). There are some case reports of spinal cord injuries in the literature, but most of them are cervical spine injuries (Bailes et al., 1991; Boden et al., 2002; Hewett et al., 2005).

We have to emphasize again that this study was only focused on moderate to severe injuries which reached the medical station after the match. In Shadgan et al. (2017) study the majority (55%) of wrestling injuries were mild, 27% were moderate and 18% were severe; no critical injury was observed. While the distribution of moderate injuries in our study differed in the three styles, the same number occurred from severe injuries. The occurrence of moderate and severe injuries was almost the same in Greco-Roman and Freestyle Wrestling, however, it was double in Women wrestling. We recorded one critical injury, which fully recovered without permanent disability.

In connection with different styles the percentage of injuries on the head and trunk was significantly higher in Freestyle Wrestling while the frequency of injuries on the extremities was significantly higher in Women wrestling. Regarding injury severity and injury location there were no significant differences between the different age groups. The analysis of the influencing factors of injury severity (moderate vs. severe & critical) showed that extremity injuries had greater odds of being severe rather than moderate compared to head injuries.

Limitation of this study: although the classification of injury severity, as defined by the UWW-MC, is precise and straightforward, but as this categorization exists only in the UWW Athena and feed-back system. Therefore, it is difficult to compare our data with other studies. There are three more aspects that limit the study: we were focusing on moderate and severe (direct and indirect) contact injuries which occurred during only matches. The documentation and the orientational diagnosis were recorded on-site based on the injury pattern followed by physical examination. We must also state that there were only five International Tournaments organized in Hungary during this period and these five were in different age groups which also complicate a direct comparison.

Conclusion

The primary purpose of this study was to investigate the specifications of contact (direct and indirect) moderate, severe and critical wrestling injuries during a series of UWW Championships by direct observation. This information can help to improve wrestling injury

prevention and enhance medical setup, preparation and coverage of wrestling competitions.

Our general injury data registered in UWW competitions correlates with the related literature considering the injuries suffered in match situations or during the competitions.

Despite the relatively low occurrence rate of injuries, there is a need for continuous education for medical teams, referees and coaches to avoid wrestling injuries (Shadgan et al., 2021). One result of this study is that a practical guideline for strangulation (Molnár Sz., 2021) was prepared as a serious preventable injury.

Acknowledgements

This work was supported by Semmelweis University under Grant EFOP-3.6.3-VEKOP-16-2017-00009. The experiments comply with the current laws of the country in which they were performed. The authors have no conflict of interest to declare. The datasets generated during and/or analyzed during the current study are not publicly available, but are available from the corresponding author who was an organizer of the study.

References

- Agel, J., Ransone, J., Dick, R., Oppliger, R. and Marshall, S.W. (2007) Descriptive epidemiology of collegiate men's wrestling injuries: National Collegiate Athletic Association Injury Surveillance System, 1988-1989 through 2003-2004. *Journal of Athletic Training* **42**, 303-10. Available from URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1941299/> [Accessed 27 Dec 2021]
- Bahr, R., Clarsen, B., Derman, W., Dvorak, J., Emery, C.A., Finch, C.F., Häggglund, M., Junge, A., Kemp, S., Khan, K.M., Marshall, S.W., Meeuwisse, W., Mountjoy, M., Orchard, J.W., Pluim, B., Quarrie, K.L., Reider, B., Schweltnus, M., Soligard, T., Stokes, K.A., Timpka, T., Verhagen, E., Bindra, A., Budgett, R., Engebretsen, L., Erdener, U. and Chamari, K. (2020) International Olympic Committee consensus statement: methods for recording and reporting of epidemiological data on injury and illness in sport 2020 (including STROBE Extension for Sport Injury and Illness Surveillance (STROBE-SIIS)). *British Journal of Sports Medicine* **54**, 372-389. <https://doi.org/10.1136/bjsports-2019-101969>
- Bailes, J.E., Hadley, M.N., Quigley, M.R., Sonntag, V.K. and Cerullo, L.J. (1991) Management of athletic injuries of the cervical spine and spinal cord. *Neurosurgery* **29**, 491-7. <https://doi.org/10.1227/00006123-199110000-00001>
- Barley, O.R., Chapman, D.W. and Abbiss, C.R. (2019) The Current State of Weight-Cutting in Combat Sports. *Sports (Basel)* **7**, 123. <https://doi.org/10.3390/sports7050123>
- Boden, B.P., Lin, W., Young, M. and Mueller, F.O. (2002) Catastrophic Injuries in Wrestlers. *The American Journal of Sports Medicine* **30**, 791-795. <https://doi.org/10.1177/03635465020300060601>
- Channon, A., Matthews, C.R. and Hillier, M. (2020) Medical care in unlicensed combat sports: A need for standardised regulatory frameworks. *Journal of Science and Medicine in Sport* **23**, 237-240. <https://doi.org/10.1016/j.jsams.2019.10.014>
- Dzsinich, C., Zsíros, L., Vallus, G., Berek, P., Nyiri, G., Pellek, S., Görög, J., Szentpétery, L. and Pataki, T. (2015) A mellkasi aorta tompa sérülése – kezelési taktika, eredmények. *Magyar Sebészet (Hungarian Journal of Surgery)* **68**, 155-166. (In Hungarian: English abstract). <https://doi.org/10.1556/1046.68.2015.4.1>
- Estwanik, J.J., Bergfeld, J. and Canty, T. (1978) Report of injuries sustained during the United States Olympic wrestling trials. *The American Journal of Sports Medicine* **6**, 335-340. <https://doi.org/10.1177/036354657800600605>
- Halloran, L. (2008) Wrestling Injuries. *Orthopaedic Nursing* **27**, 189-192. <https://doi.org/10.1097/01.NOR.0000320548.20611.16>
- Hewett, T., Pasque, C., Heyl, R. and Wroble, R. (2005) Wrestling Injuries. *Medicine and Sport Science* **48**, 152-178. <https://doi.org/10.1159/000084288>
- Jarrett, G.J., Orwin, J.F. and Dick, R.W. (1998) Injuries in Collegiate Wrestling. *The American Journal of Sports Medicine* **26**, 674-680. <https://doi.org/10.1177/03635465980260051301>
- Jlid, M., Maffulli, N., Elloumi, M., Moalla, W. and Paillard, T. (2013) Rapid weight loss alters muscular performance and perceived exertion as well as postural control in elite wrestlers. *The Journal of sports medicine and physical fitness* **53**, 620-627. Available from URL: <https://pubmed.ncbi.nlm.nih.gov/25055132/> [Accessed 27 Dec 2021]
- Kersey, R.D. and Rowan, L. (1983) Injury account during the 1980 NCAA wrestling championships. *The American Journal of Sports Medicine* **11**, 147-51. <https://doi.org/10.1177/036354658301100307>
- Khodaei, M., Olewinski, L., Shadgan, B. and Kiningham, R.R. (2015) Rapid Weight Loss in Sports with Weight Classes. *Current Sports Medicine Reports* **14**, 435-441. <https://doi.org/10.1249/JSR.0000000000000206>
- Kim, J.C. and Park, K.J. (2021) Injuries and rapid weight loss in elite Korean wrestlers: an epidemiological study. *The Physician and Sportsmedicine* **49**, 308-315, 2021/07/03. <https://doi.org/10.1080/00913847.2020.1824536>
- Kordi, R., Ziaee, V., Rostami, M. and Wallace, W.A. (2012) Sports injuries and health problems among wrestlers in Tehran. *Journal of Pakistan Medical Association* **62**, 204-8, Mar. Available from URL: https://jpma.org.pk/article-details/3291?article_id=3291 [Accessed 27 Dec 2021]
- Kroshus, E., Utter, A.C., Pierpoint, L.A., Currie, D.W., Knowles, S.B., Wasserman, E.B., Dompier, T.P., Marshall, S.W., Comstock, R.D. and Kerr, Z.Y. (2018) The First Decade of Web-Based Sports Injury Surveillance: Descriptive Epidemiology of Injuries in US High School Boys' Wrestling (2005–2006 Through 2013–2014) and National Collegiate Athletic Association Men's Wrestling (2004–2005 Through 2013–2014). *Journal of Athletic Training* **53**, 1143-1155. <https://doi.org/10.4085/1062-6050-154-17>
- Kvittem, B., Hardie, N.A., Roettger, M. and Conry, J. (1998) Incidence of Orofacial Injuries in High School Sports. *Journal of Public Health Dentistry* **58**, 288-293. <https://doi.org/10.1111/j.1752-7325.1998.tb03011.x>
- Lingor, R.J. and Olson, A. (2010) Fluid and Diet Patterns Associated with Weight Cycling and Changes in Body Composition Assessed by Continuous Monitoring Throughout a College Wrestling Season. *The Journal of Strength and Conditioning Research* **24**, 1763-1772. <https://doi.org/10.1519/JSC.0b013e3181db22fb>
- Lorish, T.R., Rizzo, T.D., Ilstrup, D.M. and Scott, S.G. (1992) Injuries in adolescent and preadolescent boys at two large wrestling tournaments. *The American Journal of Sports Medicine* **20**, 199-202. <https://doi.org/10.1177/036354659202000218>
- Maffulli, N., Longo, U.G., Gougoulias, N., Caine, D. and Denaro, V. (2010) Sport injuries: a review of outcomes. *British Medical Bulletin* **97**, 47-80. <https://doi.org/10.1093/bmb/ldq026>
- Molnár, S., Farkas, G., Rögler, G., Bacsá, P., Péteri, L. and Gáspár, K. (2014) Practical Medical Guide for Wrestling Competitions - Local, Regional or Younger Age Competitions. *International Journal of Wrestling Science* **4**, 33-36. <https://doi.org/10.1080/21615667.2014.954494>
- Molnár, S., Mensch, K. and Gáspár, K. (2020) *Wrestling*. Berlin, Heidelberg: Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-662-60752-7_86
- Molnár Sz., L.M., Nakajima K. (2021) Guidelines for the Prevention & Management of Strangulation in Olympic Wrestling. Available from URL: https://unitedworldwrestling.org/sites/default/files/2021-02/201020_strangulation_in_olympic_wrestling_report_final_version_jan2021.pdf [Accessed 07 Nov 2021].
- Myers, R.J., Linakis, S.W., Mello, M.J. and Linakis, J.G. (2010) Competitive Wrestling-related Injuries in School Aged Athletes in U.S. Emergency Departments. *The Western Journal of Emergency Medicine* **11**, 442-9. Available from URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3027436/> [Accessed 27 Dec 2021]
- Otero, J.E., Graves, C.M. and Bollier, M.J. (2017) Injuries in Collegiate Wrestlers at an Elite Division I NCAA Wrestling Program: An Epidemiological Study. *Iowa Orthopaedic Journal* **37**, 65-70. Available from URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5508281/> [Accessed 27 Dec 2021]
- Park, K.J., Lee, J.H. and Kim, H.C. (2019) Injuries in male and female elite Korean wrestling athletes: a 10-year epidemiological study. *British Journal of Sports Medicine* **53**, 430-435. <https://doi.org/10.1136/bjsports-2018-099644>

- Pasque, C.B. and Hewett, T.E. (2000) A Prospective Study of High School Wrestling Injuries. *The American Journal of Sports Medicine* **28**, 509-515.
<https://doi.org/10.1177/03635465000280041101>
- Shadgan, B., Feldman, B.J. and Jafari, S. (2010) Wrestling Injuries During the 2008 Beijing Olympic Games. *The American Journal of Sports Medicine* **38**, 1870-1876.
<https://doi.org/10.1177/0363546510369291>
- Shadgan, B., Konstantinou, L. and Molnar, S. (2021) Referees can prevent injuries in wrestling; an experience from the 2018 youth olympic games. *British Journal of Sports Medicine* **55**, A74-A74.
<http://dx.doi.org/10.1136/bjsports-2021-IOC.175>
- Shadgan, B., Molnar, S., Sikmic, S. and Chahi, A. (2017) WRESTLING INJURIES DURING THE 2016 RIO OLYMPIC GAMES. *British Journal of Sports Medicine* **51**, 387-387.
<https://doi.org/10.1136/bjsports-2016-097372.262>
- Strauss, R.H. and Lanese, R.R. (1982) Injuries among wrestlers in school and college tournaments. *Journal of the American Medical Association* **248**, 2016-9. Available from URL: <https://jamanetwork.com/journals/jama/article-abstract/378975> [Accessed 27 Dec 2021]
- Thomas, R.E. and Zamanpour, K. (2018) Injuries in wrestling: systematic review. *The Physician and Sportsmedicine* **46**, 168-196.
 doi: 10.1080/00913847.2018.1445406
- Tomin, M. and Kmetty, Á. (2021) Development of wrestling mat materials to achieve better mechanical properties and improve the safety of the athletes. *British Journal of Sports Medicine* **55**, A85-A85. <https://doi.org/10.1136/bjsports-2021-IOC.200>
- UnitedWorldWrestling. (2019a) Medical Regulations. Available from URL: https://www.org/sites/default/files/2019-07/medical_regulations_july_2019_eng_final_0.pdf [Accessed 07 Nov 2021].
- UnitedWorldWrestling. (2019b) Official Injury Report Form. Available from URL: https://unitedworldwrestling.org/sites/default/files/2019-12/injury_report_new-bs_cr_final_version_dec2019.pdf [Accessed 07 Nov 2021].
- Yamaner, F., İmamoglu, O., Atan, T., Evli, F., Karacabey, K., Sevindi, T. and Gümüşdağ, H. (2012) The Injuries of Turkish National Free-Style and Graeco-Roman Wrestlers. *Medicina Dello Sport* **65**, 549-61, 01. Available from URL: <https://www.minervamedica.it/en/journals/medicina-dello-sport/article.php?cod=R26Y2012N04A0549> [Accessed 27 Dec 2021]
- Yard, E.E., Collins, C.L., Dick, R.W. and Comstock, R.D. (2008) An Epidemiologic Comparison of High School and College Wrestling Injuries. *The American Journal of Sports Medicine* **36**, 57-64. <https://doi.org/10.1177/0363546507307507>
- Yard, E.E. and Comstock, R.D. (2008) A comparison of pediatric freestyle and Greco-Roman wrestling injuries sustained during a 2006 US national tournament. *Scandinavian Journal of Medicine & Science in Sports* **18**, 491-7.
<https://doi.org/10.1111/j.1600-0838.2007.00716.x>
- Yeo, P.C., Yeo, E.Q.Y., Probert, J., Sim, S.H.S. and Sirisena, D. (2020) A Systematic Review and Qualitative Analysis of Concussion Knowledge amongst Sports Coaches and Match Officials. *Journal of Sports Science and Medicine* **19**, 65-77.
<https://pubmed.ncbi.nlm.nih.gov/32132829/>

Key points

- We consider the UWW official classification of the injury severity to be the most straightforward with mild, moderate, severe and critical groups.
- The competition data are likely to be the most accurate because they are directly observed and recorded by medical team.
- One result of this study is that a practical guideline for strangulation was prepared as a preventable serious injury.

AUTHOR BIOGRAPHY



Szabolcs MOLNÁR

Employment

Department of Traumatology, Medical Centre Hungarian Defense Forces, Budapest, Hungary

Degree

PhD, MD

Research interests

Combat Sport Medicine and surgery

E-mail:

molnar.szabolcs.lajos@hm.gov.hu



Zsolt HUNYA

Employment

Department of Orthopedics, University of Debrecen, Debrecen, Hungary

Degree

MD

Research interests

Sport related shoulder injuries, shoulder surgery, biomechanics

E-mail:

hunya.zsolt@med.unideb.hu



Krisztián GÁSPÁR

Employment

Department of Dermatology, University of Debrecen, Hungary

Degree

PhD, MD

Research interests

Dermatological, immunological and pharmacologic aspects of combat sports medicine.

E-mail:

gaspar.krisztian@med.unideb.hu



Imre SZERB

Employment

Department of Traumatology, Semmelweis University, Budapest, Hungary

Degree


PhD, MD, med. habil

Research interests

Radiosynoviorthesis

E-mail:

szerb.imre@uzsoki.hu



Noé SZABÓ

Employment

Semmelweis University Medicine and Health Sciences, Budapest, Hungary

Degree

PhD student

Research interests

Radiosynoviorthesis

E-mail:

szabo.noe@stud.semmelweis.hu



Károly MENSCH

Employment

Department of Oral Diagnostics, Semmelweis University, Faculty of Dentistry, Budapest, Hungary

Degree

PhD

Research interests

Dentistry

E-mail:

menschkaroly@gmail.com

Éva KÖRÖSI**Employment**

Dietitian and WADA coordinator of the Hungarian Wrestling Federation, Budapest, Hungary

Degree

BSc

Research interests

Sport specific nutrition

E-mail: korosie@birkozoszov.hu

Katalin BACSKAI**Employment**

National Institute for Sports Medicine, Budapest, Hungary

Degree

MSc

Research interests

sport prevention and rehabilitation

E-mail: bacskai.katalin@osei.hu

Ákos Kálmán SÁNTHA**Employment**

Professional Medical Services and Assistance Ltd., Budapest, Hungary

Degree

MD

Research interests

Travel Medicine

E-mail:

doctor@healthguardhungary.com

Eszter Anna JANKA**Employment**

Department of Dermatology, University of Debrecen, Debrecen, Hungary

Degree

PhD

Research interests

Epidemiology and biostatistics

E-mail: janka.eszter.a@gmail.com

**Babak SHADGAN****Employment**

Department of Orthopaedics at the University of British Columbia (UBC) in Vancouver, Canada

Degree

MD, MSc Sports Med, PhD

Research interests

Injury surveillance, Sports injury prevention, Wearable and Implantable Biosensing Technologies

E-mail: shadgan@mail.ubc.ca

✉ **Szabolcs Molnár**

Department of Traumatology, Medical Centre Hungarian Defense Forces, Budapest Róbert Károly krt. 44, Hungary-1134.